

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| ALL CAUSES | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 873.1 | 728.7 | 32.9 | 18.7 | 81.6 | 108.1 | 200.0 | 431.6 | 1,004.6 | 2,428.6 | 5,688.4 | 15,321.5 |
| 1999..... | 877.0 | 731.4 | 34.7 | 19.2 | 81.2 | 108.3 | 199.2 | 427.3 | 1,021.8 | 2,484.3 | 5,751.3 | 15,476.1 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 874.7 | 799.9 | 36.5 | 21.7 | 117.5 | 149.9 | 256.9 | 552.5 | 1,253.5 | 3,015.7 | 6,854.7 | 16,605.4 |
| 1999..... | 882.0 | 801.5 | 38.5 | 22.2 | 115.9 | 150.2 | 256.7 | 546.7 | 1,280.0 | 3,109.3 | 6,999.8 | 16,931.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 871.6 | 654.3 | 29.1 | 15.6 | 44.0 | 66.7 | 143.9 | 315.8 | 777.7 | 1,945.1 | 4,900.9 | 14,768.6 |
| 1999..... | 872.2 | 658.1 | 30.8 | 16.1 | 44.7 | 66.9 | 142.5 | 313.1 | 786.5 | 1,972.9 | 4,915.4 | 14,861.2 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 915.5 | 598.4 | 29.1 | 17.5 | 75.6 | 96.2 | 180.3 | 391.8 | 948.9 | 2,375.1 | 5,652.4 | 15,532.5 |
| 1999..... | 917.7 | 596.8 | 30.7 | 17.7 | 74.5 | 96.6 | 177.6 | 386.8 | 961.5 | 2,427.8 | 5,713.2 | 15,688.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 905.8 | 656.2 | 32.5 | 20.4 | 107.6 | 134.4 | 234.3 | 502.7 | 1,177.7 | 2,949.8 | 6,817.7 | 16,897.7 |
| 1999..... | 911.2 | 658.1 | 33.9 | 20.3 | 104.9 | 134.5 | 231.5 | 494.2 | 1,200.2 | 3,043.2 | 6,965.1 | 17,202.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 924.9 | 537.9 | 25.4 | 14.4 | 41.6 | 57.7 | 125.8 | 283.2 | 735.5 | 1,893.9 | 4,860.4 | 14,948.7 |
| 1999..... | 924.1 | 532.6 | 27.4 | 14.9 | 42.2 | 58.4 | 123.3 | 281.8 | 739.1 | 1,916.2 | 4,869.7 | 15,053.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 809.6 | 1,505.6 | 56.1 | 26.2 | 120.8 | 195.5 | 360.2 | 806.1 | 1,628.6 | 3,251.5 | 6,826.4 | 14,752.1 |
| 1999..... | 817.7 | 1,551.1 | 58.9 | 28.7 | 123.1 | 191.3 | 372.3 | 808.4 | 1,683.3 | 3,343.6 | 6,883.6 | 14,787.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

ALL CAUSES

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-------|---------|------|------|-------|-------|-------|---------|---------|---------|---------|----------|
| 2000..... | 865.4 | 1,653.2 | 61.2 | 30.6 | 181.4 | 271.8 | 456.7 | 1,060.3 | 2,172.9 | 4,065.5 | 8,240.2 | 15,494.6 |
| 1999..... | 880.0 | 1,694.0 | 66.0 | 34.6 | 185.6 | 268.4 | 473.6 | 1,081.9 | 2,244.0 | 4,182.8 | 8,352.6 | 16,047.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-------|---------|------|------|------|-------|-------|-------|---------|---------|---------|----------|
| 2000..... | 759.1 | 1,352.7 | 50.8 | 21.7 | 59.6 | 126.7 | 274.7 | 594.1 | 1,213.4 | 2,659.0 | 5,967.8 | 14,441.9 |
| 1999..... | 761.3 | 1,403.3 | 51.5 | 22.6 | 60.1 | 122.0 | 282.6 | 581.0 | 1,255.7 | 2,732.1 | 6,002.7 | 14,262.9 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALEWHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

1999..... * * * * *

BLACK, FEMALE

1999..... * * * * *

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Tuberculosis (A16-A19) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.9 | 2.0 | 2.7 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 2.1 | 4.1 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.4 | 0.5 | 1.1 | 2.8 | 4.3 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.4 | 2.8 | 6.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 2.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.6 | 3.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 2.2 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.7 | 3.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.1 | 5.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.2 | 1.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.7 | * | * | * | * | * | 0.4 | 1.0 | 1.4 | 3.1 | 6.7 | * |
| 1999..... | 0.7 | * | * | * | * | * | 0.4 | 1.1 | 1.7 | 2.6 | 5.6 | 8.6 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Respiratory tuberculosis (A16) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.7 | 2.3 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 1.8 | 3.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.4 | 0.4 | 0.9 | 2.3 | 3.9 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.2 | 2.5 | 6.1 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.3 | 1.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 2.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.2 | 1.9 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.3 | 3.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.5 | 3.3 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.8 | 5.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 1.3 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | * | * | 0.4 | 0.8 | 0.9 | 2.5 | 5.3 | * |
| 1999..... | 0.6 | * | * | * | * | * | 0.4 | 0.7 | 1.5 | 2.0 | 4.8 | 7.3 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS RACE, SEX, AND YEAR | 5-14 YEARS RACE, SEX, AND YEAR | 15-24 YEARS RACE, SEX, AND YEAR | 25-34 YEARS RACE, SEX, AND YEAR | 35-44 YEARS RACE, SEX, AND YEAR | 45-54 YEARS RACE, SEX, AND YEAR | 55-64 YEARS RACE, SEX, AND YEAR | 65-74 YEARS RACE, SEX, AND YEAR | 75-84 YEARS RACE, SEX, AND YEAR | 85 YEARS AND OVER RACE, SEX, AND YEAR |
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|-----|-----|
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|-----|-----|

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|---|---|---|---|
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | * | * | * | * |
|-----------|-----|---|---|---|---|---|---|-----|---|---|---|---|

[illegible]

1999.....0.0 * * * * * 0.1 0.2 *

1999.....0.0 * * * * *

[illegible]

```
1999..... 0.1 * * * * *
```

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALE

ALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

1999..... * * * * *

BLACK, FEMALE

1999..... * * * * *

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS RACE, SEX, AND YEAR | 5-14 YEARS RACE, SEX, AND YEAR | 15-24 YEARS RACE, SEX, AND YEAR | 25-34 YEARS RACE, SEX, AND YEAR | 35-44 YEARS RACE, SEX, AND YEAR | 45-54 YEARS RACE, SEX, AND YEAR | 55-64 YEARS RACE, SEX, AND YEAR | 65-74 YEARS RACE, SEX, AND YEAR | 75-84 YEARS RACE, SEX, AND YEAR | 85 YEARS AND OVER RACE, SEX, AND YEAR |
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALEWHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Septicemia (A40-A41) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.3 | 7.1 | 0.7 | 0.2 | 0.3 | 0.7 | 2.0 | 5.0 | 12.1 | 31.4 | 80.7 | 212.9 |
| 1999..... | 11.3 | 7.4 | 0.6 | 0.2 | 0.3 | 0.7 | 1.8 | 4.7 | 11.6 | 31.6 | 79.9 | 219.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 10.1 | 7.1 | 0.7 | 0.2 | 0.3 | 0.8 | 2.0 | 5.7 | 13.2 | 34.9 | 90.3 | 211.6 |
| 1999..... | 10.1 | 9.0 | 0.6 | 0.2 | 0.3 | 0.8 | 2.1 | 5.3 | 12.7 | 35.6 | 90.9 | 224.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 12.6 | 7.2 | 0.6 | 0.1 | 0.2 | 0.6 | 1.9 | 4.2 | 11.1 | 28.5 | 74.2 | 213.4 |
| 1999..... | 12.4 | 5.8 | 0.5 | 0.2 | 0.3 | 0.7 | 1.6 | 4.2 | 10.6 | 28.3 | 72.6 | 217.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.0 | 5.6 | 0.5 | 0.2 | 0.2 | 0.6 | 1.6 | 3.9 | 10.0 | 27.8 | 74.0 | 200.7 |
| 1999..... | 10.8 | 5.4 | 0.5 | 0.2 | 0.3 | 0.6 | 1.4 | 3.7 | 9.6 | 27.6 | 73.3 | 205.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 9.6 | 5.7 | 0.6 | 0.2 | 0.2 | 0.6 | 1.7 | 4.5 | 11.0 | 31.3 | 83.2 | 200.1 |
| 1999..... | 9.5 | 6.8 | 0.5 | 0.2 | 0.3 | 0.7 | 1.6 | 4.1 | 10.1 | 31.0 | 83.7 | 210.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 12.2 | 5.5 | 0.5 | * | 0.1 | 0.5 | 1.5 | 3.3 | 9.1 | 24.9 | 67.8 | 201.0 |
| 1999..... | 12.1 | 3.9 | 0.5 | 0.2 | 0.2 | 0.5 | 1.2 | 3.3 | 9.1 | 24.8 | 66.2 | 203.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.8 | 17.3 | 1.2 | * | 0.6 | 2.1 | 4.8 | 13.9 | 31.8 | 70.0 | 176.6 | 393.6 |
| 1999..... | 16.7 | 18.8 | * | * | 0.6 | 1.6 | 5.0 | 13.4 | 30.8 | 72.9 | 171.5 | 413.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Septicemia (A40-A41) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 15.4 | 16.9 | * | * | * | 2.4 | 4.7 | 17.0 | 37.1 | 77.6 | 201.1 | 394.8 |
| 1999..... | 15.8 | 21.1 | * | * | * | 1.6 | 5.8 | 15.9 | 38.8 | 86.8 | 196.8 | 420.5 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 18.0 | 17.8 | * | * | * | 1.8 | 4.9 | 11.4 | 27.8 | 64.5 | 161.8 | 393.1 |
| 1999..... | 17.5 | 16.4 | * | * | * | 1.6 | 4.2 | 11.3 | 24.7 | 62.7 | 156.4 | 410.0 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

Syphilis (A50-A53)

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS RACE, SEX, AND YEAR | 5-14 YEARS RACE, SEX, AND YEAR | 15-24 YEARS RACE, SEX, AND YEAR | 25-34 YEARS RACE, SEX, AND YEAR | 35-44 YEARS RACE, SEX, AND YEAR | 45-54 YEARS RACE, SEX, AND YEAR | 55-64 YEARS RACE, SEX, AND YEAR | 65-74 YEARS RACE, SEX, AND YEAR | 75-84 YEARS RACE, SEX, AND YEAR | 85 YEARS AND OVER RACE, SEX, AND YEAR |
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

BLACK, FEMALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

BLACK, FEMALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Viral hepatitis (B15-B19) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | * | 0.2 | 1.7 | 5.1 | 3.8 | 4.7 | 5.4 | 3.9 |
| 1999..... | 1.8 | * | * | * | * | 0.2 | 1.6 | 4.2 | 3.4 | 4.8 | 5.4 | 4.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 2.5 | * | * | * | * | 0.2 | 2.4 | 7.8 | 5.0 | 4.8 | 6.3 | 5.4 |
| 1999..... | 2.3 | * | * | * | * | 0.3 | 2.3 | 6.5 | 4.4 | 5.2 | 6.5 | 6.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | 0.2 | 1.1 | 2.5 | 2.6 | 4.5 | 4.7 | 3.3 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.9 | 2.0 | 2.4 | 4.5 | 4.6 | 3.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | * | 0.2 | 1.7 | 4.7 | 3.2 | 4.3 | 5.2 | 3.8 |
| 1999..... | 1.7 | * | * | * | * | 0.2 | 1.5 | 3.8 | 2.9 | 4.3 | 5.1 | 4.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 2.4 | * | * | * | * | 0.2 | 2.4 | 7.3 | 4.3 | 4.4 | 6.1 | 5.4 |
| 1999..... | 2.2 | * | * | * | * | 0.3 | 2.3 | 5.8 | 4.0 | 4.9 | 6.5 | 6.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | * | * | 0.1 | 1.0 | 2.1 | 2.2 | 4.2 | 4.5 | 3.2 |
| 1999..... | 1.2 | * | * | * | * | 0.2 | 0.8 | 1.9 | 2.0 | 3.9 | 4.2 | 3.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 2.1 | 8.1 | 5.9 | 5.5 | 6.3 | * |
| 1999..... | 2.1 | * | * | * | * | * | 2.1 | 7.1 | 5.1 | 6.7 | 6.2 | 6.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Viral hepatitis (B15-B19)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|-----|-----|-----|---|
| 2000..... | 3.0 | * | * | * | * | * | 2.9 | 12.8 | 8.4 | 6.7 | 7.4 | * |
| 1999..... | 2.7 | * | * | * | * | * | 3.1 | 12.2 | 6.5 | 5.7 | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|---|
| 2000..... | 1.6 | * | * | * | * | * | 1.3 | 4.2 | 4.0 | 4.7 | 5.7 | * |
| 1999..... | 1.6 | * | * | * | * | * | 1.3 | 2.9 | 4.1 | 7.5 | 6.8 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Human immunodeficiency virus (HIV) disease (B20-B24) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.3 | * | * | 0.1 | 0.5 | 6.5 | 13.2 | 11.1 | 5.2 | 2.2 | 0.7 | * |
| 1999..... | 5.4 | * | 0.2 | 0.2 | 0.5 | 7.2 | 13.9 | 10.9 | 4.9 | 2.2 | 0.6 | * |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 8.1 | * | * | 0.1 | 0.5 | 8.6 | 19.9 | 18.1 | 8.9 | 3.8 | 1.3 | * |
| 1999..... | 8.4 | * | * | 0.2 | 0.5 | 10.2 | 21.1 | 17.9 | 8.5 | 3.9 | 1.0 | * |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.6 | * | * | 0.1 | 0.4 | 4.4 | 6.5 | 4.5 | 1.8 | 0.8 | 0.3 | * |
| 1999..... | 2.6 | * | * | 0.2 | 0.6 | 4.3 | 6.8 | 4.2 | 1.6 | 0.8 | 0.3 | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.9 | * | * | 0.1 | 0.2 | 3.3 | 7.5 | 5.9 | 2.9 | 1.0 | 0.4 | * |
| 1999..... | 3.0 | * | * | 0.1 | 0.2 | 3.9 | 8.0 | 5.8 | 2.6 | 1.1 | 0.3 | * |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 4.8 | * | * | * | 0.2 | 5.0 | 12.4 | 10.3 | 5.0 | 1.9 | 0.7 | * |
| 1999..... | 5.1 | * | * | * | 0.3 | 6.3 | 13.2 | 10.1 | 4.7 | 2.0 | 0.7 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | 1.6 | 2.6 | 1.7 | 0.8 | 0.3 | * | * |
| 1999..... | 1.0 | * | * | * | 0.2 | 1.6 | 2.7 | 1.7 | 0.6 | 0.3 | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 22.2 | * | * | 0.4 | 2.1 | 27.3 | 54.0 | 54.7 | 26.4 | 13.2 | 4.9 | * |
| 1999..... | 22.6 | * | * | 0.7 | 2.3 | 28.3 | 56.7 | 53.5 | 26.7 | 13.2 | 3.7 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Human immunodeficiency virus (HIV) disease (B20-B24)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|------|------|------|------|------|-----|---|
| 2000..... | 32.2 | * | * | * | 2.0 | 33.6 | 78.1 | 89.6 | 48.0 | 23.5 | 9.1 | * |
| 1999..... | 33.2 | * | * | 0.7 | 1.9 | 37.3 | 82.7 | 89.3 | 47.6 | 23.7 | 6.3 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|-----|------|------|------|------|-----|---|---|
| 2000..... | 13.2 | * | * | * | 2.2 | 21.7 | 32.5 | 25.5 | 9.9 | 5.7 | * | * |
| 1999..... | 13.1 | * | * | * | 2.8 | 20.2 | 33.7 | 23.8 | 10.7 | 5.6 | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.0 | 4.3 | 0.5 | 0.2 | 0.3 | 0.4 | 0.8 | 1.6 | 2.9 | 5.9 | 11.2 | 20.9 |
| 1999..... | 2.7 | 5.3 | 0.5 | 0.2 | 0.2 | 0.6 | 1.2 | 2.3 | 3.8 | 7.9 | 14.3 | 26.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 2.1 | 4.8 | 0.5 | 0.2 | 0.3 | 0.4 | 1.0 | 2.0 | 3.3 | 7.3 | 13.4 | 26.2 |
| 1999..... | 2.7 | 6.1 | 0.7 | 0.2 | 0.3 | 0.7 | 1.5 | 2.7 | 4.3 | 9.3 | 17.0 | 30.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | 3.9 | 0.4 | 0.2 | 0.2 | 0.4 | 0.6 | 1.2 | 2.6 | 4.8 | 9.7 | 18.7 |
| 1999..... | 2.6 | 4.5 | 0.4 | 0.2 | 0.2 | 0.6 | 0.9 | 2.0 | 3.4 | 6.8 | 12.5 | 25.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.0 | 3.3 | 0.4 | 0.2 | 0.2 | 0.3 | 0.7 | 1.4 | 2.7 | 5.8 | 11.0 | 20.8 |
| 1999..... | 2.6 | 4.5 | 0.5 | 0.2 | 0.2 | 0.5 | 1.0 | 1.9 | 3.4 | 7.6 | 13.8 | 26.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 2.1 | 3.8 | 0.4 | 0.2 | 0.3 | 0.3 | 0.8 | 1.8 | 3.1 | 7.1 | 13.1 | 25.6 |
| 1999..... | 2.6 | 5.5 | 0.6 | 0.2 | 0.2 | 0.5 | 1.3 | 2.3 | 3.9 | 8.9 | 16.1 | 29.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | 2.8 | 0.4 | 0.2 | 0.2 | 0.2 | 0.5 | 1.1 | 2.3 | 4.7 | 9.7 | 18.7 |
| 1999..... | 2.5 | 3.4 | * | 0.2 | 0.2 | 0.4 | 0.6 | 1.5 | 3.0 | 6.5 | 12.2 | 25.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.3 | 10.5 | * | * | 0.4 | 1.0 | 1.7 | 3.1 | 4.9 | 7.6 | 13.6 | 24.1 |
| 1999..... | 3.6 | 10.5 | 1.0 | * | 0.3 | 1.7 | 3.0 | 5.6 | 7.3 | 11.6 | 20.9 | 34.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|------|---|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 2.6 | 10.5 | * | * | * | 0.9 | 2.2 | 4.0 | 5.9 | 9.8 | 17.6 | 29.8 |
| 1999..... | 3.8 | 10.7 | * | * | * | 1.9 | 3.1 | 6.4 | 8.2 | 13.4 | 28.2 | 42.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|---|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 2.1 | 10.5 | * | * | * | 1.0 | 1.2 | 2.3 | 4.2 | 6.0 | 11.1 | 21.8 |
| 1999..... | 3.5 | 10.4 | * | * | * | 1.5 | 3.0 | 4.9 | 6.6 | 10.2 | 16.6 | 30.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms (C00-C97) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 200.9 | 2.4 | 2.8 | 2.6 | 4.5 | 10.5 | 36.8 | 129.3 | 371.3 | 826.4 | 1,340.8 | 1,795.6 |
| 1999..... | 201.6 | 1.8 | 2.8 | 2.6 | 4.6 | 10.6 | 37.3 | 130.4 | 380.8 | 836.2 | 1,340.0 | 1,796.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 212.5 | 2.5 | 3.0 | 2.8 | 5.2 | 10.0 | 32.9 | 133.3 | 423.5 | 1,014.0 | 1,730.9 | 2,571.9 |
| 1999..... | 214.5 | 1.8 | 2.9 | 2.7 | 5.3 | 10.4 | 33.6 | 135.1 | 437.9 | 1,031.2 | 1,746.2 | 2,619.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 189.9 | 2.2 | 2.5 | 2.3 | 3.7 | 11.0 | 40.6 | 125.5 | 323.6 | 671.8 | 1,077.4 | 1,461.3 |
| 1999..... | 189.4 | 1.7 | 2.6 | 2.4 | 3.8 | 10.7 | 41.1 | 125.9 | 328.7 | 676.6 | 1,068.1 | 1,449.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 212.2 | 2.5 | 2.7 | 2.6 | 4.4 | 10.3 | 34.6 | 122.9 | 362.8 | 822.6 | 1,333.8 | 1,795.6 |
| 1999..... | 212.5 | 1.8 | 2.9 | 2.5 | 4.4 | 10.3 | 35.3 | 123.9 | 370.4 | 831.5 | 1,331.5 | 1,795.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 222.5 | 2.7 | 3.1 | 2.9 | 5.1 | 10.0 | 31.0 | 125.0 | 406.8 | 999.3 | 1,707.1 | 2,569.2 |
| 1999..... | 224.2 | 2.0 | 3.2 | 2.7 | 5.1 | 10.2 | 32.0 | 126.0 | 419.9 | 1,017.1 | 1,723.2 | 2,612.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 202.2 | 2.2 | 2.4 | 2.4 | 3.6 | 10.6 | 38.2 | 120.9 | 321.7 | 674.7 | 1,080.1 | 1,464.7 |
| 1999..... | 201.1 | 1.5 | 2.7 | 2.3 | 3.7 | 10.5 | 38.7 | 121.9 | 324.3 | 677.3 | 1,067.6 | 1,453.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 175.5 | * | 3.1 | 2.4 | 5.2 | 12.4 | 54.5 | 197.3 | 502.4 | 980.0 | 1,595.5 | 2,004.0 |
| 1999..... | 177.7 | * | 2.2 | 2.9 | 5.5 | 12.7 | 54.5 | 199.3 | 530.2 | 998.8 | 1,609.8 | 2,012.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms (C00-C97)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-------|---|-----|-----|-----|------|------|-------|-------|---------|---------|---------|
| 2000..... | 195.6 | * | 3.0 | 2.7 | 6.1 | 10.5 | 48.8 | 223.2 | 654.3 | 1,303.5 | 2,283.6 | 3,012.7 |
| 1999..... | 198.4 | * | 2.3 | 3.0 | 6.4 | 11.9 | 47.7 | 232.3 | 689.2 | 1,316.4 | 2,302.5 | 3,098.1 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-------|---|-----|-----|-----|------|------|-------|-------|-------|---------|---------|
| 2000..... | 157.2 | * | 3.3 | 2.0 | 4.2 | 14.1 | 59.5 | 175.7 | 386.6 | 744.5 | 1,177.6 | 1,582.6 |
| 1999..... | 159.0 | * | 2.1 | 2.8 | 4.7 | 13.4 | 60.5 | 171.9 | 408.9 | 767.3 | 1,194.4 | 1,560.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.7 | * | * | * | 0.1 | 0.1 | 0.7 | 2.9 | 6.8 | 10.8 | 13.2 | 19.2 |
| 1999..... | 2.7 | * | * | * | 0.1 | 0.1 | 0.6 | 3.0 | 6.9 | 10.4 | 14.4 | 19.2 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 3.7 | * | * | * | * | 0.2 | 1.0 | 4.5 | 11.1 | 16.2 | 18.9 | 25.7 |
| 1999..... | 3.7 | * | * | * | 0.1 | 0.1 | 0.9 | 4.7 | 11.0 | 16.0 | 20.7 | 26.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.8 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.0 | 6.4 | 9.3 | 16.4 |
| 1999..... | 1.8 | * | * | * | * | * | 0.4 | 1.3 | 3.1 | 5.9 | 10.1 | 16.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.7 | * | * | * | * | 0.1 | 0.5 | 2.4 | 6.1 | 10.4 | 13.2 | 19.5 |
| 1999..... | 2.7 | * | * | * | * | 0.1 | 0.5 | 2.4 | 6.2 | 10.0 | 14.2 | 19.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 3.5 | * | * | * | * | 0.1 | 0.7 | 3.7 | 9.7 | 15.2 | 18.4 | 26.3 |
| 1999..... | 3.6 | * | * | * | * | * | 0.7 | 3.7 | 9.8 | 15.3 | 20.1 | 26.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | * | * | 0.3 | 1.2 | 2.7 | 6.3 | 9.6 | 16.6 |
| 1999..... | 1.9 | * | * | * | * | * | 0.3 | 1.1 | 2.9 | 5.7 | 10.3 | 16.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.1 | * | * | * | * | * | 1.1 | 6.1 | 13.7 | 15.1 | 15.3 | 15.4 |
| 1999..... | 3.3 | * | * | * | * | * | 1.2 | 7.4 | 13.7 | 14.9 | 17.2 | 14.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|------|------|------|------|
| 2000..... | 5.0 | * | * | * | * | * | 1.8 | 11.0 | 25.4 | 27.0 | 27.9 | * |
| 1999..... | 5.0 | * | * | * | * | * | 1.8 | 12.7 | 24.2 | 24.2 | 30.9 | 22.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|------|
| 2000..... | 1.5 | * | * | * | * | * | * | 2.0 | 4.8 | 6.4 | 7.7 | 14.7 |
| 1999..... | 1.7 | * | * | * | * | * | 0.7 | 2.9 | 5.7 | 8.1 | 9.0 | 10.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of esophagus (C15) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | * | 0.1 | 0.7 | 3.6 | 10.9 | 20.2 | 25.7 | 25.6 |
| 1999..... | 4.4 | * | * | * | * | 0.1 | 0.7 | 3.5 | 10.8 | 20.3 | 24.6 | 26.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.9 | * | * | * | * | 0.2 | 1.2 | 6.4 | 19.0 | 34.5 | 45.0 | 45.3 |
| 1999..... | 6.8 | * | * | * | * | 0.2 | 1.1 | 6.3 | 18.7 | 35.7 | 43.9 | 46.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.1 | * | * | * | * | * | 0.1 | 1.0 | 3.5 | 8.4 | 12.7 | 17.1 |
| 1999..... | 2.0 | * | * | * | * | * | 0.2 | 0.9 | 3.7 | 7.7 | 11.7 | 18.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | * | * | * | * | 0.1 | 0.7 | 3.4 | 10.1 | 19.8 | 25.4 | 26.3 |
| 1999..... | 4.5 | * | * | * | * | 0.1 | 0.6 | 3.2 | 10.0 | 19.7 | 24.3 | 27.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 7.2 | * | * | * | * | 0.2 | 1.2 | 6.0 | 17.8 | 34.1 | 44.9 | 47.1 |
| 1999..... | 7.0 | * | * | * | * | 0.2 | 1.1 | 5.8 | 17.3 | 34.9 | 43.5 | 46.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.1 | * | * | * | * | * | 0.1 | 0.8 | 3.0 | 7.9 | 12.1 | 17.4 |
| 1999..... | 2.0 | * | * | * | * | * | 0.1 | 0.7 | 3.2 | 7.0 | 11.4 | 19.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.7 | * | * | * | * | * | 0.8 | 6.8 | 20.0 | 27.5 | 34.3 | 22.3 |
| 1999..... | 4.7 | * | * | * | * | * | 1.0 | 6.7 | 20.3 | 29.3 | 31.7 | 24.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of esophagus (C15)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|------|------|------|------|
| 2000..... | 7.0 | * | * | * | * | * | 1.2 | 11.7 | 35.5 | 44.9 | 56.7 | 35.1 |
| 1999..... | 7.3 | * | * | * | * | * | 1.7 | 11.9 | 36.4 | 48.6 | 55.5 | 47.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|------|------|------|
| 2000..... | 2.6 | * | * | * | * | * | * | 2.7 | 8.2 | 14.8 | 20.7 | 16.9 |
| 1999..... | 2.4 | * | * | * | * | * | * | 2.5 | 8.1 | 15.1 | 17.5 | 14.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of stomach (C16) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | * | * | * | * | 0.3 | 1.1 | 3.0 | 7.5 | 16.8 | 31.1 | 52.0 |
| 1999..... | 4.7 | * | * | * | * | 0.4 | 1.1 | 3.1 | 8.0 | 17.3 | 31.8 | 49.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 5.4 | * | * | * | * | 0.4 | 1.3 | 3.9 | 10.7 | 24.0 | 43.2 | 71.8 |
| 1999..... | 5.6 | * | * | * | * | 0.4 | 1.3 | 4.3 | 11.2 | 24.5 | 46.0 | 68.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.8 | * | * | * | * | 0.3 | 0.9 | 2.1 | 4.6 | 10.9 | 23.0 | 43.5 |
| 1999..... | 3.8 | * | * | * | * | 0.3 | 1.0 | 1.9 | 5.0 | 11.4 | 22.3 | 42.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.3 | * | * | * | * | 0.3 | 0.9 | 2.5 | 6.6 | 14.7 | 27.3 | 46.8 |
| 1999..... | 4.4 | * | * | * | * | 0.3 | 0.9 | 2.5 | 6.7 | 15.4 | 28.3 | 44.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 5.1 | * | * | * | * | 0.3 | 1.0 | 3.2 | 9.4 | 21.2 | 38.3 | 65.7 |
| 1999..... | 5.2 | * | * | * | * | 0.4 | 1.0 | 3.6 | 9.5 | 22.0 | 41.0 | 61.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.5 | * | * | * | * | 0.2 | 0.7 | 1.7 | 3.9 | 9.3 | 19.9 | 38.7 |
| 1999..... | 3.5 | * | * | * | * | 0.2 | 0.8 | 1.5 | 4.1 | 9.9 | 19.6 | 37.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.2 | * | * | * | * | 0.7 | 2.1 | 5.8 | 14.0 | 31.2 | 65.1 | 101.9 |
| 1999..... | 6.3 | * | * | * | * | 0.7 | 2.2 | 6.1 | 16.1 | 30.5 | 65.8 | 102.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of stomach (C16) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 7.3 | * | * | * | * | 0.8 | 2.9 | 8.3 | 21.6 | 44.6 | 88.5 | 125.6 |
| 1999..... | 7.6 | * | * | * | * | * | 2.7 | 8.9 | 24.1 | 44.2 | 98.1 | 136.9 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 5.2 | * | * | * | * | * | 1.3 | 3.8 | 8.3 | 21.4 | 50.9 | 92.0 |
| 1999..... | 5.1 | * | * | * | * | * | 1.8 | 3.8 | 10.0 | 20.5 | 46.4 | 88.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of colon, rectum and anus (C18-C21) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 20.9 | * | * | * | 0.1 | 0.7 | 3.0 | 11.4 | 33.1 | 77.6 | 146.6 | 266.8 |
| 1999..... | 21.0 | * | * | * | 0.1 | 0.7 | 3.0 | 11.3 | 34.1 | 78.2 | 147.6 | 270.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 21.1 | * | * | * | 0.1 | 0.8 | 3.4 | 13.0 | 40.3 | 96.6 | 176.5 | 299.5 |
| 1999..... | 21.2 | * | * | * | 0.1 | 0.8 | 3.0 | 12.9 | 41.4 | 96.1 | 179.7 | 311.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 20.6 | * | * | * | * | 0.6 | 2.7 | 9.9 | 26.6 | 62.0 | 126.4 | 252.8 |
| 1999..... | 20.7 | * | * | * | 0.1 | 0.6 | 2.9 | 9.9 | 27.3 | 63.5 | 126.1 | 252.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 21.9 | * | * | * | 0.1 | 0.7 | 2.8 | 10.5 | 31.5 | 76.1 | 144.4 | 266.2 |
| 1999..... | 22.0 | * | * | * | 0.1 | 0.7 | 2.8 | 10.6 | 32.5 | 76.6 | 145.4 | 270.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 22.0 | * | * | * | * | 0.8 | 3.1 | 12.1 | 38.1 | 94.8 | 174.0 | 298.8 |
| 1999..... | 22.2 | * | * | * | * | 0.8 | 2.9 | 12.1 | 39.4 | 95.1 | 177.4 | 314.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 21.7 | * | * | * | * | 0.6 | 2.4 | 9.0 | 25.3 | 60.5 | 124.2 | 252.3 |
| 1999..... | 21.7 | * | * | * | * | 0.6 | 2.6 | 9.1 | 26.0 | 61.2 | 123.8 | 251.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 19.5 | * | * | * | * | 1.0 | 5.1 | 19.9 | 50.8 | 102.0 | 193.6 | 310.8 |
| 1999..... | 19.4 | * | * | * | * | 1.1 | 4.8 | 18.3 | 53.6 | 104.0 | 196.8 | 306.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of colon, rectum and anus (C18-C21)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|------|------|-------|-------|-------|
| 2000..... | 19.8 | * | * | * | * | 1.0 | 6.0 | 22.9 | 65.3 | 125.7 | 234.0 | 348.0 |
| 1999..... | 19.3 | * | * | * | * | 1.2 | 4.7 | 21.1 | 68.8 | 121.7 | 238.5 | 324.9 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|------|------|------|-------|-------|
| 2000..... | 19.1 | * | * | * | * | 1.0 | 4.4 | 17.4 | 39.8 | 84.8 | 169.1 | 295.3 |
| 1999..... | 19.6 | * | * | * | * | 0.9 | 4.9 | 15.9 | 41.9 | 91.1 | 171.8 | 299.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of liver and intrahepatic bile ducts (C22) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.7 | * | 0.2 | 0.1 | 0.1 | 0.2 | 1.0 | 4.5 | 9.4 | 18.9 | 28.9 | 32.3 |
| 1999..... | 4.5 | * | 0.2 | * | 0.1 | 0.2 | 1.0 | 4.2 | 8.9 | 19.2 | 28.2 | 30.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.1 | * | * | * | 0.1 | 0.3 | 1.3 | 7.3 | 14.1 | 27.4 | 41.3 | 42.4 |
| 1999..... | 5.8 | * | 0.3 | * | 0.1 | 0.3 | 1.4 | 6.6 | 13.1 | 27.9 | 38.5 | 38.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.4 | * | * | * | * | 0.1 | 0.6 | 1.9 | 5.2 | 12.0 | 20.5 | 27.9 |
| 1999..... | 3.4 | * | * | * | * | 0.2 | 0.6 | 1.8 | 5.1 | 12.0 | 21.3 | 27.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | * | * | * | 0.1 | 0.1 | 0.7 | 3.7 | 8.2 | 17.5 | 27.9 | 31.6 |
| 1999..... | 4.4 | * | 0.2 | * | 0.1 | 0.1 | 0.8 | 3.5 | 7.7 | 18.0 | 27.3 | 29.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 5.8 | * | * | * | * | 0.2 | 1.0 | 5.8 | 11.9 | 25.3 | 40.4 | 41.1 |
| 1999..... | 5.6 | * | * | * | * | 0.1 | 1.1 | 5.3 | 11.3 | 26.2 | 37.6 | 39.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.4 | * | * | * | * | * | 0.4 | 1.7 | 4.7 | 11.0 | 19.3 | 27.6 |
| 1999..... | 3.4 | * | * | * | * | * | 0.5 | 1.6 | 4.4 | 11.2 | 20.5 | 25.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.5 | * | * | * | * | 0.4 | 1.8 | 8.8 | 14.3 | 23.0 | 29.2 | 31.7 |
| 1999..... | 4.3 | * | * | * | * | 0.4 | 1.6 | 7.5 | 13.4 | 23.7 | 29.9 | 34.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|------|------|------|------|
| 2000..... | 6.3 | * | * | * | * | * | 2.5 | 15.4 | 24.5 | 35.7 | 38.5 | 44.7 |
| 1999..... | 5.7 | * | * | * | * | * | 2.4 | 13.5 | 20.9 | 35.1 | 40.2 | 28.3 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|------|------|------|
| 2000..... | 2.9 | * | * | * | * | * | 1.2 | 3.3 | 6.4 | 13.8 | 23.6 | 26.2 |
| 1999..... | 3.1 | * | * | * | * | * | 0.9 | 2.5 | 7.7 | 15.4 | 23.8 | 37.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of pancreas (C25) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 10.7 | * | * | * | * | 0.1 | 1.2 | 6.4 | 19.6 | 44.6 | 76.8 | 95.1 |
| 1999..... | 10.7 | * | * | * | * | 0.1 | 1.3 | 6.6 | 20.1 | 45.5 | 73.5 | 99.2 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 10.6 | * | * | * | * | 0.2 | 1.5 | 8.2 | 24.0 | 51.1 | 84.4 | 97.6 |
| 1999..... | 10.6 | * | * | * | * | 0.2 | 1.7 | 8.5 | 24.4 | 53.0 | 80.9 | 103.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.7 | * | * | * | * | 0.1 | 1.0 | 4.6 | 15.6 | 39.3 | 71.7 | 94.1 |
| 1999..... | 10.7 | * | * | * | * | 0.1 | 1.0 | 4.8 | 16.2 | 39.4 | 68.6 | 97.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.2 | * | * | * | * | 0.2 | 1.2 | 6.2 | 19.0 | 43.7 | 75.5 | 93.6 |
| 1999..... | 11.2 | * | * | * | * | 0.1 | 1.3 | 6.3 | 19.4 | 44.7 | 72.2 | 98.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 11.1 | * | * | * | * | 0.2 | 1.4 | 7.9 | 23.3 | 50.6 | 83.8 | 97.6 |
| 1999..... | 11.1 | * | * | * | * | 0.2 | 1.6 | 8.1 | 23.3 | 52.4 | 79.9 | 104.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 11.2 | * | * | * | * | * | 0.9 | 4.4 | 15.1 | 38.0 | 69.9 | 91.9 |
| 1999..... | 11.2 | * | * | * | * | * | 0.9 | 4.6 | 15.7 | 38.4 | 66.9 | 96.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 9.7 | * | * | * | * | * | 1.7 | 9.3 | 27.9 | 57.6 | 99.7 | 120.1 |
| 1999..... | 9.7 | * | * | * | * | * | 2.0 | 9.9 | 28.7 | 58.3 | 95.8 | 118.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of pancreas (C25) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 9.3 | * | * | * | * | * | 2.0 | 12.3 | 35.7 | 62.9 | 103.8 | 101.1 |
| 1999..... | 9.7 | * | * | * | * | * | 2.4 | 13.8 | 38.1 | 65.3 | 104.1 | 106.5 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.0 | * | * | * | * | * | 1.4 | 6.7 | 21.9 | 53.7 | 97.1 | 128.1 |
| 1999..... | 9.7 | * | * | * | * | * | 1.6 | 6.7 | 21.5 | 53.2 | 90.9 | 124.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of larynx (C32) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | * | 0.1 | 1.1 | 3.6 | 6.7 | 8.1 | 6.4 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 1.1 | 3.9 | 6.8 | 7.3 | 7.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 2.3 | * | * | * | * | * | 0.2 | 1.9 | 6.3 | 11.5 | 15.4 | 16.5 |
| 1999..... | 2.2 | * | * | * | * | * | 0.3 | 2.0 | 6.6 | 11.6 | 14.0 | 17.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.2 | 2.8 | 3.1 | 2.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.4 | 2.9 | 2.9 | 2.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | * | 0.1 | 1.0 | 3.1 | 6.2 | 7.7 | 6.5 |
| 1999..... | 1.4 | * | * | * | * | * | 0.1 | 0.9 | 3.4 | 6.5 | 7.2 | 6.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | * | * | * | 0.2 | 1.5 | 5.3 | 10.5 | 14.8 | 16.8 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 1.6 | 5.8 | 10.9 | 13.6 | 16.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.4 | 1.1 | 2.6 | 2.9 | 2.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.2 | 2.8 | 2.9 | 2.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | * | * | * | * | 3.0 | 9.1 | 12.4 | 14.3 | 6.9 |
| 1999..... | 1.9 | * | * | * | * | * | 0.6 | 3.0 | 8.7 | 11.3 | 10.2 | 8.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of larynx (C32)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|------|
| 2000..... | 3.4 | * | * | * | * | * | * | 5.3 | 17.9 | 23.2 | 27.6 | * |
| 1999..... | 3.2 | * | * | * | * | * | 0.9 | 5.4 | 16.7 | 20.9 | 22.2 | 22.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|-----|---|
| 2000..... | 0.8 | * | * | * | * | * | * | 1.0 | 2.3 | 4.6 | 6.3 | * |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.9 | 2.7 | 4.3 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of trachea, bronchus and lung (C33-C34) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 56.5 | * | * | * | 0.1 | 0.5 | 6.1 | 32.1 | 124.0 | 287.6 | 372.2 | 298.1 |
| 1999..... | 55.8 | * | * | * | 0.1 | 0.5 | 6.1 | 31.9 | 125.5 | 284.6 | 364.4 | 295.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 67.2 | * | * | * | * | 0.5 | 6.9 | 39.2 | 156.9 | 382.5 | 523.2 | 494.5 |
| 1999..... | 67.1 | * | * | * | * | 0.5 | 6.6 | 39.2 | 161.5 | 381.6 | 521.4 | 509.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 46.3 | * | * | * | * | 0.5 | 5.4 | 25.2 | 94.0 | 209.5 | 270.3 | 213.5 |
| 1999..... | 45.0 | * | * | * | * | 0.4 | 5.5 | 25.0 | 92.6 | 205.2 | 259.3 | 205.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 60.5 | * | * | * | * | 0.5 | 5.7 | 30.5 | 123.7 | 291.1 | 375.7 | 301.1 |
| 1999..... | 59.5 | * | * | * | * | 0.5 | 5.8 | 30.2 | 124.6 | 287.2 | 366.0 | 298.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 70.8 | * | * | * | * | 0.5 | 6.1 | 36.1 | 152.6 | 380.6 | 521.1 | 498.3 |
| 1999..... | 70.6 | * | * | * | * | 0.5 | 6.1 | 35.9 | 156.5 | 378.9 | 518.0 | 515.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 50.6 | * | * | * | * | 0.5 | 5.2 | 24.9 | 96.7 | 216.2 | 276.9 | 216.8 |
| 1999..... | 48.9 | * | * | * | * | 0.4 | 5.4 | 24.7 | 94.9 | 211.0 | 263.6 | 207.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 45.5 | * | * | * | * | 0.6 | 10.2 | 51.6 | 154.5 | 307.9 | 382.0 | 288.8 |
| 1999..... | 45.8 | * | * | * | * | 0.4 | 9.4 | 52.3 | 160.8 | 306.7 | 395.4 | 291.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|---|------|------|-------|-------|-------|-------|
| 2000..... | 60.5 | * | * | * | * | * | 13.5 | 73.7 | 233.5 | 467.2 | 623.9 | 519.3 |
| 1999..... | 61.2 | * | * | * | * | * | 11.4 | 76.2 | 248.5 | 465.6 | 645.1 | 532.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|---|-----|------|------|-------|-------|-------|
| 2000..... | 31.9 | * | * | * | * | * | 7.3 | 33.2 | 94.2 | 191.9 | 235.0 | 192.5 |
| 1999..... | 31.8 | * | * | * | * | * | 7.7 | 32.5 | 93.9 | 190.9 | 245.6 | 190.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant melanoma of skin (C43) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.7 | * | * | * | 0.1 | 0.5 | 1.4 | 3.2 | 5.4 | 9.0 | 13.2 | 18.7 |
| 1999..... | 2.6 | * | * | * | 0.1 | 0.5 | 1.4 | 3.1 | 5.4 | 9.1 | 13.5 | 16.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 3.4 | * | * | * | 0.1 | 0.6 | 1.6 | 4.2 | 7.6 | 13.0 | 20.4 | 30.2 |
| 1999..... | 3.4 | * | * | * | 0.1 | 0.6 | 1.8 | 4.1 | 7.6 | 13.4 | 20.7 | 26.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | * | 0.1 | 0.4 | 1.2 | 2.3 | 3.5 | 5.7 | 8.3 | 13.8 |
| 1999..... | 1.9 | * | * | * | * | 0.4 | 1.0 | 2.2 | 3.4 | 5.5 | 8.7 | 12.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.2 | * | * | * | 0.1 | 0.6 | 1.6 | 3.8 | 6.2 | 10.1 | 14.3 | 20.1 |
| 1999..... | 3.2 | * | * | * | 0.1 | 0.6 | 1.7 | 3.7 | 6.2 | 10.2 | 14.6 | 17.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 4.1 | * | * | * | 0.1 | 0.7 | 1.9 | 4.9 | 8.6 | 14.5 | 22.2 | 33.1 |
| 1999..... | 4.0 | * | * | * | 0.1 | 0.7 | 2.1 | 4.7 | 8.6 | 15.0 | 22.4 | 28.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.4 | * | * | * | * | 0.5 | 1.4 | 2.6 | 4.0 | 6.4 | 9.0 | 14.5 |
| 1999..... | 2.3 | * | * | * | * | 0.5 | 1.2 | 2.6 | 3.9 | 6.2 | 9.4 | 12.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.8 | 1.8 | 3.6 | 6.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 1.5 | 2.8 | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of breast (C50) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 15.4 | * | * | * | * | 1.2 | 6.3 | 17.2 | 31.6 | 49.7 | 79.2 | 145.5 |
| 1999..... | 15.2 | * | * | * | * | 1.2 | 6.1 | 17.2 | 31.6 | 50.0 | 79.5 | 143.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.2 | 3.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.0 | 4.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 29.8 | * | * | * | * | 2.5 | 12.5 | 33.4 | 59.7 | 89.4 | 131.2 | 206.4 |
| 1999..... | 29.5 | * | * | * | * | 2.3 | 12.1 | 33.5 | 59.9 | 89.9 | 131.3 | 202.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.0 | * | * | * | * | 1.1 | 5.7 | 16.0 | 30.5 | 49.9 | 79.6 | 145.6 |
| 1999..... | 15.8 | * | * | * | * | 1.0 | 5.4 | 16.0 | 30.4 | 49.6 | 79.5 | 145.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.2 | 3.7 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 1.9 | 4.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 31.1 | * | * | * | * | 2.2 | 11.4 | 31.4 | 58.3 | 90.5 | 132.2 | 206.3 |
| 1999..... | 30.7 | * | * | * | * | 2.0 | 10.8 | 31.4 | 58.1 | 89.9 | 131.7 | 205.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 15.4 | * | * | * | * | 2.3 | 11.3 | 28.5 | 46.0 | 57.1 | 92.2 | 176.9 |
| 1999..... | 15.6 | * | * | * | * | 2.4 | 11.8 | 29.7 | 47.0 | 62.0 | 96.3 | 147.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of breast (C50) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 28.9 | * | * | * | * | 4.4 | 21.1 | 52.0 | 80.0 | 97.5 | 146.4 | 247.2 |
| 1999..... | 29.3 | * | * | * | * | 4.5 | 21.9 | 54.0 | 81.7 | 105.5 | 152.0 | 205.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of cervix uteri (C53) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | 0.6 | 1.5 | 2.5 | 3.1 | 3.8 | 4.9 | 7.2 |
| 1999..... | 1.5 | * | * | * | * | 0.6 | 1.6 | 2.6 | 3.3 | 3.8 | 4.7 | 7.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.0 | * | * | * | * | 1.2 | 3.1 | 4.8 | 6.0 | 6.9 | 8.1 | 10.3 |
| 1999..... | 3.0 | * | * | * | * | 1.1 | 3.1 | 5.0 | 6.3 | 6.9 | 7.8 | 11.2 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | 0.6 | 1.4 | 2.2 | 2.7 | 3.1 | 4.3 | 6.2 |
| 1999..... | 1.4 | * | * | * | * | 0.6 | 1.4 | 2.4 | 2.8 | 3.2 | 4.1 | 6.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.7 | * | * | * | * | 1.2 | 2.8 | 4.3 | 5.2 | 5.8 | 7.2 | 8.9 |
| 1999..... | 2.8 | * | * | * | * | 1.2 | 2.9 | 4.8 | 5.5 | 5.8 | 6.8 | 9.6 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.5 | * | * | * | * | 1.0 | 2.6 | 4.4 | 6.3 | 9.3 | 12.1 | 19.8 |
| 1999..... | 2.4 | * | * | * | * | 0.7 | 2.5 | 3.6 | 7.2 | 10.5 | 11.5 | 21.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.7 | * | * | * | * | 1.9 | 5.0 | 8.0 | 11.2 | 16.1 | 19.5 | 28.0 |
| 1999..... | 4.6 | * | * | * | * | 1.3 | 4.8 | 6.5 | 12.7 | 18.1 | 18.4 | 30.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

ALL AGES 1/
UNDER 1 YEAR 2/
1-4 YEARS
5-14 YEARS
15-24 YEARS
25-34 YEARS
35-44 YEARS
45-54 YEARS
55-64 YEARS
65-74 YEARS
75-84 YEARS
85 YEARS
AND OVER

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|-----|------|------|
| 2000..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.7 | 9.8 | 15.9 | 24.4 |
| 1999..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.4 | 9.9 | 16.5 | 23.9 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 4.7 | * | * | * | * | 0.1 | 0.6 | 2.7 | 8.9 | 17.9 | 26.6 | 34.9 |
| 1999..... | 4.6 | * | * | * | * | 0.1 | 0.6 | 2.6 | 8.3 | 18.0 | 27.6 | 34.0 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|-----|------|------|
| 2000..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.3 | 8.7 | 14.9 | 24.1 |
| 1999..... | 2.4 | * | * | * | * | * | 0.3 | 1.3 | 4.1 | 9.0 | 15.9 | 23.3 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 4.7 | * | * | * | * | 0.1 | 0.6 | 2.7 | 8.3 | 15.9 | 25.0 | 34.4 |
| 1999..... | 4.7 | * | * | * | * | * | 0.6 | 2.5 | 7.8 | 16.5 | 26.5 | 33.1 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|------|------|------|
| 2000..... | 3.0 | * | * | * | * | * | 0.4 | 1.6 | 8.4 | 22.1 | 31.1 | 33.2 |
| 1999..... | 2.8 | * | * | * | * | * | 0.5 | 1.5 | 8.1 | 20.9 | 27.8 | 35.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|------|
| 2000..... | 5.7 | * | * | * | * | * | 0.7 | 3.0 | 14.8 | 38.2 | 50.0 | 47.1 |
| 1999..... | 5.4 | * | * | * | * | * | 0.9 | 2.7 | 14.2 | 36.0 | 44.4 | 50.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of ovary (C56) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.1 | * | * | * | 0.1 | 0.3 | 1.1 | 4.3 | 10.2 | 19.8 | 33.3 | 40.4 |
| 1999..... | 5.0 | * | * | * | 0.1 | 0.2 | 1.1 | 4.3 | 10.4 | 19.8 | 31.9 | 37.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.0 | * | * | * | 0.1 | 0.5 | 2.1 | 8.3 | 19.5 | 36.1 | 55.8 | 57.8 |
| 1999..... | 9.8 | * | * | * | 0.1 | 0.4 | 2.2 | 8.5 | 19.8 | 36.0 | 53.3 | 53.2 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.6 | * | * | * | * | 0.3 | 1.1 | 4.4 | 10.6 | 20.4 | 34.3 | 41.5 |
| 1999..... | 5.5 | * | * | * | 0.1 | 0.2 | 1.1 | 4.5 | 10.7 | 20.4 | 32.6 | 38.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 11.1 | * | * | * | * | 0.6 | 2.2 | 8.7 | 20.5 | 37.5 | 57.7 | 59.3 |
| 1999..... | 10.8 | * | * | * | 0.2 | 0.4 | 2.3 | 8.8 | 20.8 | 37.3 | 54.5 | 54.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.0 | * | * | * | * | * | 1.1 | 3.4 | 8.5 | 17.6 | 27.1 | 34.5 |
| 1999..... | 3.1 | * | * | * | * | * | 0.8 | 3.7 | 9.3 | 18.1 | 29.2 | 28.8 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 5.8 | * | * | * | * | * | 2.0 | 6.2 | 15.0 | 30.4 | 43.6 | 48.9 |
| 1999..... | 5.9 | * | * | * | * | * | 1.6 | 6.7 | 16.3 | 31.3 | 46.8 | 40.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of prostate (C61) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.3 | * | * | * | * | * | 0.1 | 1.0 | 8.0 | 37.5 | 106.2 | 206.6 |
| 1999..... | 11.6 | * | * | * | * | * | 0.1 | 1.0 | 8.3 | 40.2 | 110.6 | 206.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 23.1 | * | * | * | * | * | 0.1 | 2.0 | 16.9 | 83.0 | 263.4 | 686.1 |
| 1999..... | 23.8 | * | * | * | * | * | 0.1 | 2.1 | 17.5 | 89.2 | 275.8 | 695.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.2 | * | * | * | * | * | * | 0.8 | 6.8 | 33.0 | 97.3 | 196.4 |
| 1999..... | 11.6 | * | * | * | * | * | 0.1 | 0.9 | 7.0 | 35.8 | 102.5 | 194.8 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 22.8 | * | * | * | * | * | * | 1.7 | 14.1 | 72.3 | 240.4 | 655.7 |
| 1999..... | 23.6 | * | * | * | * | * | 0.1 | 1.8 | 14.5 | 78.9 | 254.6 | 659.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 15.1 | * | * | * | * | * | * | 2.5 | 20.6 | 87.8 | 234.1 | 357.5 |
| 1999..... | 15.4 | * | * | * | * | * | * | 2.7 | 22.0 | 91.4 | 230.1 | 367.1 |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 31.9 | * | * | * | * | * | * | 5.4 | 47.6 | 208.5 | 619.5 | 1,213.2 |
| 1999..... | 32.4 | * | * | * | * | * | * | 5.9 | 50.8 | 216.8 | 613.9 | 1,248.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of kidney and renal pelvis (C64-C65) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.3 | * | * | 0.1 | 0.1 | 0.1 | 0.7 | 3.3 | 9.2 | 17.3 | 26.7 | 33.3 |
| 1999..... | 4.1 | * | * | 0.1 | * | 0.1 | 0.8 | 3.2 | 9.0 | 16.9 | 24.8 | 31.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 5.4 | * | * | * | * | 0.1 | 1.0 | 4.8 | 13.4 | 24.5 | 37.6 | 52.3 |
| 1999..... | 5.2 | * | * | * | * | 0.2 | 1.1 | 4.7 | 13.2 | 23.9 | 35.2 | 47.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.2 | * | * | * | * | * | 0.5 | 1.9 | 5.3 | 11.4 | 19.3 | 25.2 |
| 1999..... | 3.0 | * | * | * | * | * | 0.5 | 1.8 | 5.2 | 11.2 | 17.8 | 24.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | * | * | * | * | 0.1 | 0.7 | 3.4 | 9.4 | 17.7 | 27.0 | 34.1 |
| 1999..... | 4.4 | * | * | 0.1 | * | 0.1 | 0.7 | 3.2 | 9.3 | 17.3 | 25.1 | 31.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 5.8 | * | * | * | * | * | 1.0 | 4.9 | 13.6 | 24.8 | 38.0 | 53.9 |
| 1999..... | 5.5 | * | * | * | * | 0.1 | 1.0 | 4.6 | 13.5 | 24.3 | 36.0 | 48.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.5 | * | * | * | * | * | 0.4 | 1.9 | 5.4 | 11.8 | 19.5 | 25.6 |
| 1999..... | 3.3 | * | * | * | * | * | 0.4 | 1.8 | 5.4 | 11.5 | 17.8 | 24.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.0 | * | * | * | * | * | 0.9 | 3.5 | 8.9 | 16.1 | 26.1 | 29.8 |
| 1999..... | 3.0 | * | * | * | * | 0.4 | 1.1 | 3.9 | 8.8 | 15.8 | 24.0 | 31.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of kidney and renal pelvis (C64-C65)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|------|
| 2000..... | 3.8 | * | * | * | * | * | 1.0 | 5.1 | 13.7 | 25.2 | 37.0 | 45.8 |
| 1999..... | 3.8 | * | * | * | * | * | 1.6 | 6.1 | 13.8 | 22.6 | 31.2 | 45.6 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|------|------|------|
| 2000..... | 2.2 | * | * | * | * | * | 0.8 | 2.2 | 5.2 | 9.5 | 19.5 | 23.1 |
| 1999..... | 2.3 | * | * | * | * | * | * | 2.1 | 5.0 | 10.8 | 19.6 | 25.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of bladder (C67) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | * | * | 0.2 | 1.2 | 5.1 | 15.7 | 35.8 | 68.7 |
| 1999..... | 4.4 | * | * | * | * | * | 0.2 | 1.3 | 5.3 | 16.0 | 35.5 | 68.1 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.1 | * | * | * | * | * | 0.3 | 1.8 | 8.0 | 25.5 | 61.5 | 132.2 |
| 1999..... | 6.1 | * | * | * | * | * | 0.3 | 2.0 | 8.4 | 26.0 | 61.8 | 129.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.7 | * | * | * | * | * | 0.2 | 0.6 | 2.4 | 7.6 | 18.4 | 41.4 |
| 1999..... | 2.7 | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 7.9 | 18.0 | 42.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.9 | * | * | * | * | * | 0.3 | 1.2 | 5.4 | 16.4 | 36.5 | 71.2 |
| 1999..... | 4.8 | * | * | * | * | * | 0.2 | 1.3 | 5.4 | 16.6 | 36.6 | 69.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 6.9 | * | * | * | * | * | 0.3 | 1.9 | 8.5 | 26.9 | 63.7 | 138.9 |
| 1999..... | 6.8 | * | * | * | * | * | 0.3 | 2.0 | 8.7 | 27.3 | 64.1 | 136.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.0 | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 7.6 | 18.1 | 42.3 |
| 1999..... | 2.9 | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 7.6 | 18.0 | 41.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.4 | * | * | * | * | * | * | 1.1 | 4.1 | 13.0 | 35.3 | 49.9 |
| 1999..... | 2.6 | * | * | * | * | * | * | 1.6 | 5.7 | 14.7 | 29.7 | 60.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of bladder (C67)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|------|------|------|
| 2000..... | 2.8 | * | * | * | * | * | * | 1.4 | 6.0 | 16.8 | 51.2 | 81.9 |
| 1999..... | 3.0 | * | * | * | * | * | * | 2.7 | 7.7 | 18.4 | 45.0 | 80.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|------|------|------|
| 2000..... | 2.1 | * | * | * | * | * | * | * | 2.7 | 10.2 | 25.7 | 36.5 |
| 1999..... | 2.3 | * | * | * | * | * | * | * | 4.1 | 12.0 | 20.5 | 52.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | 0.5 | 0.7 | 0.8 | 0.6 | 1.1 | 2.4 | 5.2 | 9.7 | 16.7 | 20.1 | 16.7 |
| 1999..... | 4.7 | * | 0.7 | 0.8 | 0.6 | 1.2 | 2.4 | 5.3 | 10.5 | 17.1 | 20.5 | 15.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 5.2 | * | 0.7 | 0.9 | 0.7 | 1.4 | 3.2 | 6.3 | 12.1 | 20.5 | 23.9 | 21.3 |
| 1999..... | 5.3 | * | 0.7 | 0.8 | 0.7 | 1.5 | 2.9 | 6.4 | 12.7 | 21.1 | 25.2 | 19.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.0 | * | 0.6 | 0.7 | 0.5 | 0.8 | 1.7 | 4.1 | 7.6 | 13.6 | 17.6 | 14.7 |
| 1999..... | 4.1 | * | 0.8 | 0.8 | 0.5 | 0.9 | 1.9 | 4.2 | 8.5 | 13.9 | 17.4 | 14.2 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.2 | * | 0.7 | 0.8 | 0.6 | 1.2 | 2.6 | 5.7 | 10.5 | 18.0 | 21.1 | 17.3 |
| 1999..... | 5.3 | * | 0.8 | 0.8 | 0.6 | 1.3 | 2.6 | 5.7 | 11.3 | 18.5 | 21.5 | 16.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 5.8 | * | 0.7 | 0.9 | 0.7 | 1.5 | 3.4 | 6.8 | 12.9 | 22.0 | 25.1 | 22.5 |
| 1999..... | 5.9 | * | 0.8 | 0.8 | 0.7 | 1.6 | 3.1 | 6.8 | 13.6 | 22.7 | 26.5 | 21.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.5 | * | 0.6 | 0.7 | 0.5 | 0.9 | 1.8 | 4.5 | 8.2 | 14.7 | 18.3 | 15.1 |
| 1999..... | 4.7 | * | 0.8 | 0.8 | 0.5 | 1.0 | 2.1 | 4.6 | 9.2 | 15.0 | 18.2 | 14.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | 0.8 | 0.6 | 0.7 | 1.6 | 3.0 | 5.8 | 7.9 | 12.1 | 10.7 |
| 1999..... | 2.2 | * | * | 0.9 | 0.6 | 0.7 | 1.4 | 3.3 | 6.1 | 8.0 | 12.0 | 9.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|------|---|
| 2000..... | 2.5 | * | * | 1.0 | 0.7 | 1.1 | 2.0 | 3.9 | 7.4 | 9.4 | 12.6 | * |
| 1999..... | 2.4 | * | * | 0.8 | 0.8 | 0.9 | 1.8 | 4.6 | 7.5 | 9.6 | 12.3 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|---|---|-----|-----|-----|-----|------|------|
| 2000..... | 1.9 | * | * | 0.7 | * | * | 1.2 | 2.2 | 4.5 | 6.8 | 11.8 | 11.1 |
| 1999..... | 2.0 | * | * | 1.0 | * | * | 1.0 | 2.2 | 5.0 | 6.9 | 11.9 | 9.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 20.5 | 1.0 | 1.0 | 0.9 | 1.9 | 2.5 | 4.1 | 10.8 | 30.4 | 77.3 | 149.8 | 201.9 |
| 1999..... | 20.7 | 0.8 | 0.9 | 0.9 | 1.9 | 2.5 | 4.3 | 11.0 | 32.3 | 78.7 | 148.9 | 198.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 22.2 | 1.1 | 1.0 | 1.1 | 2.3 | 2.9 | 4.8 | 13.2 | 36.8 | 98.0 | 189.4 | 267.4 |
| 1999..... | 22.3 | * | 1.0 | 1.0 | 2.3 | 3.0 | 5.1 | 13.5 | 39.7 | 98.4 | 186.8 | 268.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 18.9 | * | 0.9 | 0.8 | 1.4 | 2.0 | 3.4 | 8.6 | 24.6 | 60.2 | 123.1 | 173.7 |
| 1999..... | 19.1 | * | 0.8 | 0.8 | 1.5 | 2.1 | 3.4 | 8.6 | 25.6 | 62.5 | 123.6 | 168.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 22.3 | 1.2 | 1.0 | 1.0 | 1.8 | 2.4 | 4.0 | 10.5 | 30.4 | 79.1 | 153.7 | 206.3 |
| 1999..... | 22.3 | 0.8 | 1.0 | 0.9 | 1.9 | 2.5 | 4.2 | 10.8 | 32.0 | 80.4 | 152.2 | 203.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 24.1 | * | 1.0 | 1.1 | 2.2 | 2.8 | 4.8 | 12.8 | 36.6 | 100.2 | 194.9 | 276.8 |
| 1999..... | 24.1 | * | 1.0 | 1.0 | 2.3 | 2.8 | 5.0 | 13.3 | 39.3 | 100.6 | 191.3 | 277.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 20.5 | * | 1.0 | 0.8 | 1.3 | 1.9 | 3.2 | 8.3 | 24.5 | 61.5 | 125.7 | 176.2 |
| 1999..... | 20.7 | * | 0.9 | 0.7 | 1.5 | 2.1 | 3.3 | 8.5 | 25.2 | 63.6 | 125.8 | 173.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 14.4 | * | 0.9 | 0.7 | 2.3 | 3.1 | 5.4 | 15.1 | 36.3 | 73.1 | 123.2 | 175.9 |
| 1999..... | 14.6 | * | * | 1.0 | 2.3 | 3.0 | 5.6 | 14.3 | 39.5 | 74.7 | 129.2 | 162.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 15.2 | * | * | 0.8 | 2.7 | 3.7 | 5.7 | 18.7 | 46.5 | 91.7 | 150.5 | 207.5 |
| 1999..... | 15.6 | * | * | 1.2 | 2.6 | 3.9 | 6.9 | 18.2 | 49.6 | 91.7 | 157.2 | 204.3 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 13.6 | * | * | * | 1.8 | 2.5 | 5.1 | 12.1 | 28.5 | 59.5 | 106.6 | 162.7 |
| 1999..... | 13.6 | * | * | 0.8 | 2.1 | 2.2 | 4.4 | 11.0 | 31.9 | 62.2 | 112.5 | 144.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Hodgkin's disease (C81) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.7 | 1.3 | 2.1 | 2.4 |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.8 | 1.6 | 2.2 | 2.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.8 | 1.6 | 2.4 | 3.5 |
| 1999..... | 0.6 | * | * | * | 0.3 | 0.5 | 0.5 | 0.6 | 1.0 | 2.0 | 2.4 | 3.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.6 | 1.0 | 1.8 | 2.0 |
| 1999..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 | 1.2 | 2.0 | 1.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.5 | 0.7 | 1.3 | 2.2 | 2.6 |
| 1999..... | 0.6 | * | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.8 | 1.6 | 2.3 | 2.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.6 | * | * | * | 0.2 | 0.5 | 0.4 | 0.6 | 0.7 | 1.7 | 2.6 | 3.7 |
| 1999..... | 0.6 | * | * | * | 0.3 | 0.5 | 0.5 | 0.6 | 1.1 | 2.0 | 2.6 | 3.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.6 | 1.0 | 2.0 | 2.1 |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 | 1.3 | 2.1 | 1.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | 0.4 | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | 0.5 | * | * | * | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Non-Hodgkin's lymphoma (C82-C85) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 8.3 | * | * | 0.1 | 0.5 | 0.8 | 1.7 | 4.7 | 13.2 | 30.8 | 61.2 | 79.2 |
| 1999..... | 8.4 | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 4.9 | 13.5 | 32.1 | 62.2 | 77.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 8.8 | * | * | 0.1 | 0.6 | 0.9 | 2.2 | 6.1 | 16.2 | 37.8 | 73.2 | 100.4 |
| 1999..... | 8.8 | * | * | 0.1 | 0.5 | 0.9 | 2.3 | 6.3 | 16.6 | 38.7 | 73.8 | 100.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 7.8 | * | * | * | 0.3 | 0.6 | 1.2 | 3.3 | 10.5 | 25.0 | 53.1 | 70.0 |
| 1999..... | 7.9 | * | * | * | 0.3 | 0.6 | 1.2 | 3.4 | 10.7 | 26.7 | 54.4 | 67.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 9.2 | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 4.7 | 13.8 | 32.5 | 64.4 | 83.4 |
| 1999..... | 9.3 | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 5.0 | 14.0 | 33.9 | 65.3 | 81.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 9.7 | * | * | 0.2 | 0.6 | 0.9 | 2.2 | 6.1 | 16.7 | 39.7 | 77.1 | 106.7 |
| 1999..... | 9.8 | * | * | 0.1 | 0.5 | 0.9 | 2.2 | 6.4 | 16.9 | 40.8 | 77.1 | 104.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 8.7 | * | * | * | 0.2 | 0.6 | 1.1 | 3.4 | 11.2 | 26.5 | 55.8 | 73.4 |
| 1999..... | 8.9 | * | * | * | 0.3 | 0.6 | 1.2 | 3.5 | 11.3 | 28.3 | 57.3 | 71.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.9 | * | * | * | 0.6 | 1.1 | 2.1 | 5.1 | 10.7 | 18.9 | 27.3 | 36.4 |
| 1999..... | 4.1 | * | * | * | 0.6 | 1.1 | 2.4 | 4.9 | 11.3 | 19.7 | 32.4 | 40.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Non-Hodgkin's lymphoma (C82-C85) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 4.3 | * | * | * | * | 1.2 | 2.6 | 7.2 | 15.1 | 23.8 | 30.3 | 40.4 |
| 1999..... | 4.7 | * | * | * | * | 1.6 | 3.3 | 6.9 | 15.3 | 24.6 | 39.6 | 55.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.5 | * | * | * | * | 0.9 | 1.7 | 3.3 | 7.2 | 15.3 | 25.5 | 34.7 |
| 1999..... | 3.6 | * | * | * | * | * | 1.7 | 3.2 | 8.2 | 16.2 | 28.1 | 34.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Leukemia (C91-C95) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 7.8 | 1.0 | 0.9 | 0.8 | 1.2 | 1.3 | 1.8 | 3.8 | 10.1 | 27.6 | 54.7 | 82.5 |
| 1999..... | 7.7 | 0.7 | 0.8 | 0.8 | 1.3 | 1.3 | 1.8 | 3.8 | 10.8 | 27.5 | 53.2 | 82.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 8.7 | 1.1 | 0.9 | 0.9 | 1.5 | 1.5 | 1.9 | 4.4 | 12.7 | 37.7 | 74.4 | 113.7 |
| 1999..... | 8.6 | * | 0.8 | 0.9 | 1.5 | 1.5 | 2.0 | 4.4 | 13.4 | 37.2 | 72.4 | 114.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 6.8 | * | 0.9 | 0.6 | 0.9 | 1.0 | 1.7 | 3.3 | 7.7 | 19.3 | 41.3 | 69.0 |
| 1999..... | 6.8 | * | 0.7 | 0.7 | 1.0 | 1.2 | 1.6 | 3.2 | 8.4 | 19.5 | 40.3 | 68.2 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 8.5 | 1.2 | 1.0 | 0.8 | 1.2 | 1.2 | 1.8 | 3.7 | 10.1 | 28.7 | 56.7 | 85.1 |
| 1999..... | 8.4 | 0.8 | 0.9 | 0.8 | 1.3 | 1.3 | 1.8 | 3.8 | 10.9 | 28.4 | 54.8 | 85.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 9.5 | * | 0.9 | 1.0 | 1.4 | 1.4 | 1.9 | 4.2 | 12.7 | 39.1 | 77.2 | 117.9 |
| 1999..... | 9.4 | * | 0.9 | 0.8 | 1.5 | 1.4 | 2.0 | 4.3 | 13.5 | 38.4 | 74.7 | 120.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 7.5 | * | 1.0 | 0.7 | 0.9 | 0.9 | 1.6 | 3.2 | 7.7 | 20.1 | 42.8 | 71.1 |
| 1999..... | 7.4 | * | 0.8 | 0.7 | 1.0 | 1.2 | 1.6 | 3.2 | 8.5 | 20.0 | 41.3 | 70.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.9 | * | * | 0.6 | 1.4 | 1.7 | 2.1 | 4.9 | 10.8 | 22.1 | 38.9 | 66.2 |
| 1999..... | 4.9 | * | * | 0.9 | 1.4 | 1.3 | 1.9 | 4.5 | 10.7 | 24.2 | 42.0 | 55.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Leukemia (C91-C95) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 5.5 | * | * | 0.7 | 1.9 | 2.0 | 2.0 | 5.7 | 14.5 | 30.2 | 53.2 | 90.5 |
| 1999..... | 5.4 | * | * | 1.0 | 1.6 | 1.5 | 2.0 | 5.8 | 14.0 | 31.7 | 53.4 | 71.7 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | 0.9 | 1.4 | 2.1 | 4.3 | 8.0 | 16.2 | 30.2 | 56.0 |
| 1999..... | 4.5 | * | * | 0.7 | 1.3 | 1.1 | 1.8 | 3.4 | 8.3 | 18.8 | 35.1 | 49.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Multiple myeloma and immunoproliferative neoplasms (C88,C90) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.0 | * | * | * | * | * | 0.3 | 1.9 | 6.4 | 17.5 | 31.6 | 37.3 |
| 1999..... | 4.0 | * | * | * | * | 0.1 | 0.3 | 1.9 | 7.2 | 17.4 | 31.2 | 36.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 4.1 | * | * | * | * | * | 0.4 | 2.2 | 7.1 | 20.7 | 38.9 | 49.1 |
| 1999..... | 4.2 | * | * | * | * | * | 0.4 | 2.2 | 8.7 | 20.4 | 37.9 | 49.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.9 | * | * | * | * | * | 0.3 | 1.6 | 5.8 | 14.9 | 26.7 | 32.3 |
| 1999..... | 3.9 | * | * | * | * | * | 0.3 | 1.6 | 5.9 | 15.0 | 26.7 | 30.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.0 | * | * | * | * | * | 0.3 | 1.6 | 5.6 | 16.4 | 30.1 | 34.8 |
| 1999..... | 4.0 | * | * | * | * | * | 0.3 | 1.6 | 6.2 | 16.3 | 29.6 | 34.8 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 4.2 | * | * | * | * | * | 0.3 | 1.8 | 6.4 | 19.5 | 37.6 | 47.9 |
| 1999..... | 4.2 | * | * | * | * | * | 0.3 | 1.9 | 7.7 | 19.2 | 36.5 | 48.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.8 | * | * | * | * | * | 0.2 | 1.4 | 5.0 | 13.8 | 25.0 | 29.2 |
| 1999..... | 3.8 | * | * | * | * | * | 0.2 | 1.2 | 4.8 | 13.9 | 25.0 | 29.2 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.2 | * | * | * | * | * | 0.8 | 4.6 | 14.1 | 31.3 | 56.1 | 71.8 |
| 1999..... | 5.2 | * | * | * | * | * | 0.9 | 4.4 | 16.9 | 29.6 | 54.1 | 63.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|------|
| 2000..... | 5.0 | * | * | * | * | * | * | 5.2 | 15.7 | 36.8 | 65.9 | 73.4 |
| 1999..... | 5.0 | * | * | * | * | * | 1.1 | 4.6 | 19.4 | 33.8 | 63.0 | 75.0 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|------|
| 2000..... | 5.4 | * | * | * | * | * | 0.9 | 4.1 | 12.8 | 27.3 | 50.2 | 71.1 |
| 1999..... | 5.3 | * | * | * | * | * | 0.7 | 4.2 | 15.0 | 26.6 | 48.8 | 58.4 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS RACE, SEX, AND YEAR | 5-14 YEARS RACE, SEX, AND YEAR | 15-24 YEARS RACE, SEX, AND YEAR | 25-34 YEARS RACE, SEX, AND YEAR | 35-44 YEARS RACE, SEX, AND YEAR | 45-54 YEARS RACE, SEX, AND YEAR | 55-64 YEARS RACE, SEX, AND YEAR | 65-74 YEARS RACE, SEX, AND YEAR | 75-84 YEARS RACE, SEX, AND YEAR | 85 YEARS AND OVER RACE, SEX, AND YEAR |
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|------|-----|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 22.9 | 0.6 | 0.9 | 0.7 | 1.3 | 1.9 | 4.7 | 14.3 | 38.8 | 86.5 | 153.2 | 236.6 |
| 1999..... | 24.3 | * | 0.8 | 0.7 | 1.4 | 2.0 | 5.2 | 15.3 | 42.2 | 92.4 | 161.8 | 249.2 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|------|---|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 2000..... | 23.6 | * | 1.0 | 0.7 | 1.6 | 2.3 | 5.1 | 16.3 | 45.7 | 104.5 | 184.6 | 285.3 |
| 1999..... | 25.1 | * | 0.8 | 0.8 | 1.7 | 2.6 | 5.8 | 17.7 | 50.2 | 111.7 | 193.8 | 306.9 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 22.3 | * | 0.7 | 0.6 | 1.0 | 1.4 | 4.2 | 12.4 | 32.5 | 71.7 | 132.0 | 215.7 |
| 1999..... | 23.5 | * | 0.7 | 0.7 | 1.0 | 1.4 | 4.5 | 13.0 | 34.9 | 76.6 | 140.4 | 224.8 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|------|---|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 24.3 | * | 0.8 | 0.7 | 1.3 | 1.9 | 4.3 | 13.4 | 37.8 | 86.5 | 153.7 | 241.0 |
| 1999..... | 25.7 | * | 0.8 | 0.7 | 1.4 | 2.0 | 4.9 | 14.5 | 40.8 | 92.5 | 162.2 | 251.9 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|------|---|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 2000..... | 24.9 | * | 1.0 | 0.7 | 1.6 | 2.5 | 4.7 | 15.3 | 43.8 | 104.7 | 185.1 | 292.8 |
| 1999..... | 26.5 | * | 0.9 | 0.8 | 1.6 | 2.6 | 5.6 | 16.5 | 48.3 | 111.5 | 195.3 | 313.6 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|-----|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 23.8 | * | 0.6 | 0.6 | 1.1 | 1.4 | 3.9 | 11.6 | 32.3 | 71.3 | 132.3 | 218.8 |
| 1999..... | 25.0 | * | 0.7 | 0.6 | 1.1 | 1.4 | 4.2 | 12.6 | 33.8 | 76.7 | 139.8 | 226.1 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|------|---|-----|-----|-----|-----|-----|------|------|-------|-------|-------|
| 2000..... | 19.0 | * | 1.1 | 0.7 | 1.5 | 1.7 | 7.6 | 22.6 | 52.5 | 97.6 | 165.0 | 211.4 |
| 1999..... | 20.3 | * | * | 0.8 | 1.5 | 2.0 | 7.5 | 23.7 | 60.6 | 102.5 | 174.2 | 243.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|-----|-----|------|------|-------|-------|-------|
| 2000..... | 20.1 | * | * | 0.8 | 1.9 | 1.9 | 8.6 | 27.3 | 71.1 | 117.2 | 204.3 | 234.1 |
| 1999..... | 21.4 | * | * | 0.7 | 2.0 | 2.4 | 8.4 | 30.4 | 79.6 | 128.1 | 201.3 | 271.7 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 17.9 | * | * | * | 1.0 | 1.5 | 6.7 | 18.7 | 38.4 | 83.4 | 141.1 | 201.9 |
| 1999..... | 19.2 | * | * | 0.9 | 0.9 | 1.7 | 6.8 | 18.1 | 46.1 | 83.9 | 158.0 | 231.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

ALL AGES 1/
UNDER 1 YEAR 2/
1-4 YEARS
5-14 YEARS
15-24 YEARS
25-34 YEARS
35-44 YEARS
45-54 YEARS
55-64 YEARS
65-74 YEARS
75-84 YEARS
85 YEARS AND OVER

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| 2000..... | 4.9 | 1.6 | 0.3 | 0.2 | 0.2 | 0.4 | 0.8 | 1.9 | 5.1 | 14.0 | 37.7 | 83.8 |
| 1999..... | 4.9 | 1.5 | 0.4 | 0.3 | 0.3 | 0.5 | 0.8 | 1.8 | 4.8 | 14.7 | 36.8 | 82.5 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 4.9 | 1.6 | 0.4 | 0.2 | 0.3 | 0.4 | 0.8 | 2.1 | 5.7 | 17.2 | 46.7 | 103.3 |
| 1999..... | 4.9 | 1.5 | 0.4 | 0.2 | 0.3 | 0.5 | 0.9 | 2.1 | 5.9 | 18.6 | 45.5 | 105.1 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| 2000..... | 5.0 | 1.5 | 0.3 | 0.3 | 0.2 | 0.4 | 0.8 | 1.6 | 4.5 | 11.4 | 31.6 | 75.4 |
| 1999..... | 4.8 | 1.5 | 0.4 | 0.3 | 0.2 | 0.4 | 0.6 | 1.6 | 3.9 | 11.6 | 31.0 | 72.9 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| 2000..... | 5.4 | 1.6 | 0.4 | 0.2 | 0.2 | 0.4 | 0.8 | 1.8 | 5.1 | 14.1 | 38.7 | 86.4 |
| 1999..... | 5.3 | 1.4 | 0.4 | 0.3 | 0.2 | 0.4 | 0.7 | 1.8 | 4.8 | 14.9 | 37.6 | 85.2 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 5.3 | 1.6 | 0.4 | 0.2 | 0.3 | 0.5 | 0.8 | 2.1 | 5.7 | 17.2 | 48.2 | 107.4 |
| 1999..... | 5.3 | 1.4 | 0.4 | 0.3 | 0.2 | 0.4 | 0.9 | 2.0 | 5.9 | 18.9 | 46.7 | 110.8 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| 2000..... | 5.5 | 1.6 | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.5 | 4.6 | 11.5 | 32.3 | 77.4 |
| 1999..... | 5.2 | 1.4 | 0.4 | 0.3 | 0.2 | 0.4 | 0.6 | 1.5 | 3.8 | 11.6 | 31.4 | 74.4 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|------|------|------|
| 2000..... | 3.2 | * | * | * | * | 0.6 | 1.1 | 2.8 | 5.6 | 14.6 | 31.6 | 64.3 |
| 1999..... | 3.4 | * | * | * | 0.5 | 0.6 | 1.2 | 2.7 | 6.4 | 15.4 | 33.6 | 62.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 3.1 | * | * | * | * | * | 1.2 | 2.9 | 7.3 | 18.3 | 35.0 | 70.2 |
| 1999..... | 3.1 | * | * | * | * | 0.8 | 1.1 | 2.6 | 7.4 | 18.2 | 37.5 | 58.7 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 3.3 | * | * | * | * | 0.8 | 1.0 | 2.6 | 4.3 | 11.9 | 29.5 | 61.8 |
| 1999..... | 3.6 | * | * | * | * | * | 1.2 | 2.8 | 5.6 | 13.3 | 31.3 | 63.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Anemias (D50-D64) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.6 | * | 0.2 | 0.1 | 0.2 | 0.4 | 0.4 | 0.6 | 1.1 | 2.8 | 9.5 | 41.8 |
| 1999..... | 1.7 | * | 0.2 | 0.1 | 0.3 | 0.4 | 0.4 | 0.6 | 1.0 | 2.7 | 9.6 | 43.6 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | 0.1 | 0.2 | 0.4 | 0.3 | 0.6 | 1.1 | 2.9 | 9.4 | 38.0 |
| 1999..... | 1.3 | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.7 | 0.9 | 2.8 | 9.9 | 41.8 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | 0.1 | 0.2 | 0.4 | 0.5 | 0.6 | 1.0 | 2.8 | 9.6 | 43.4 |
| 1999..... | 2.0 | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.6 | 1.0 | 2.6 | 9.3 | 44.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.8 | 2.6 | 9.3 | 41.6 |
| 1999..... | 1.5 | * | * | 0.1 | 0.1 | * | 0.1 | 0.3 | 0.6 | 2.4 | 9.4 | 42.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 9.1 | 37.3 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 2.5 | 9.8 | 42.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 9.4 | 43.4 |
| 1999..... | 1.9 | * | * | * | 0.1 | * | 0.1 | 0.3 | 0.7 | 2.4 | 9.1 | 43.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.7 | * | * | 0.5 | 1.1 | 2.4 | 2.5 | 3.1 | 3.4 | 5.2 | 13.6 | 51.1 |
| 1999..... | 2.9 | * | 0.9 | 0.5 | 1.5 | 2.6 | 2.5 | 3.2 | 4.3 | 4.9 | 13.6 | 53.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Anemias (D50-D64)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|-----|------|------|
| 2000..... | 2.4 | * | * | * | 1.0 | 2.2 | 2.1 | 3.4 | 3.5 | 4.3 | 13.5 | 51.1 |
| 1999..... | 2.8 | * | * | * | 1.7 | 2.9 | 2.8 | 3.4 | 5.3 | 5.2 | 12.6 | 41.3 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|-----|------|------|
| 2000..... | 3.1 | * | * | * | 1.1 | 2.5 | 2.8 | 2.8 | 3.4 | 5.9 | 13.6 | 51.1 |
| 1999..... | 3.1 | * | * | * | 1.2 | 2.4 | 2.3 | 3.0 | 3.5 | 4.6 | 14.2 | 57.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Diabetes mellitus (E10-E14) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 25.2 | * | * | 0.1 | 0.4 | 1.7 | 4.3 | 13.3 | 38.3 | 91.8 | 180.2 | 315.6 |
| 1999..... | 25.1 | * | * | 0.1 | 0.4 | 1.5 | 4.3 | 13.2 | 38.9 | 92.8 | 179.1 | 315.6 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 23.5 | * | * | * | 0.5 | 2.0 | 5.2 | 15.5 | 43.2 | 100.3 | 196.2 | 327.4 |
| 1999..... | 23.4 | * | * | * | 0.4 | 1.7 | 5.1 | 15.9 | 43.5 | 101.0 | 195.8 | 332.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 26.8 | * | * | * | 0.4 | 1.3 | 3.4 | 11.2 | 33.8 | 84.8 | 169.3 | 310.5 |
| 1999..... | 26.7 | * | * | * | 0.4 | 1.4 | 3.6 | 10.6 | 34.7 | 86.1 | 167.9 | 308.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 24.6 | * | * | * | 0.3 | 1.4 | 3.8 | 11.2 | 33.4 | 82.4 | 166.4 | 300.1 |
| 1999..... | 24.3 | * | * | 0.1 | 0.4 | 1.3 | 3.7 | 11.1 | 32.8 | 82.9 | 166.1 | 298.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 23.4 | * | * | * | 0.4 | 1.8 | 4.7 | 13.4 | 38.4 | 93.2 | 187.4 | 321.8 |
| 1999..... | 23.2 | * | * | * | 0.4 | 1.5 | 4.5 | 13.7 | 37.9 | 93.7 | 187.9 | 323.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 25.7 | * | * | * | 0.3 | 1.0 | 3.0 | 9.1 | 28.7 | 73.3 | 152.1 | 290.8 |
| 1999..... | 25.4 | * | * | * | 0.3 | 1.2 | 3.0 | 8.5 | 28.1 | 73.9 | 151.5 | 288.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 34.1 | * | * | * | 0.9 | 3.6 | 8.1 | 31.0 | 83.5 | 183.7 | 357.2 | 526.3 |
| 1999..... | 34.2 | * | * | * | 0.8 | 3.1 | 9.0 | 31.3 | 91.7 | 186.9 | 339.9 | 530.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Diabetes mellitus (E10-E14)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|-----|-----|------|------|------|-------|-------|-------|
| 2000..... | 28.4 | * | * | * | 1.0 | 4.0 | 9.5 | 34.8 | 92.2 | 176.8 | 327.2 | 417.2 |
| 1999..... | 28.7 | * | * | * | * | 3.4 | 10.4 | 36.4 | 98.1 | 179.2 | 307.2 | 469.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|-----|-----|-----|------|------|-------|-------|-------|
| 2000..... | 39.1 | * | * | * | 0.9 | 3.3 | 6.9 | 27.8 | 77.0 | 188.7 | 375.4 | 571.8 |
| 1999..... | 39.2 | * | * | * | 0.9 | 2.7 | 7.7 | 27.1 | 86.9 | 192.5 | 359.5 | 555.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Nutritional deficiencies (E40-E64) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.5 | 10.4 | 49.4 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.4 | 10.3 | 53.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.3 | 0.8 | 2.7 | 9.5 | 41.4 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.7 | 10.1 | 46.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.4 | 11.1 | 52.8 |
| 1999..... | 2.0 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.1 | 10.4 | 56.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.3 | 10.0 | 48.3 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.0 | 9.6 | 52.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 9.0 | 40.9 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 9.2 | 45.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.1 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.3 | 10.7 | 51.5 |
| 1999..... | 2.1 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.9 | 9.8 | 56.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | * | * | 0.6 | 1.7 | 4.6 | 16.4 | 67.1 |
| 1999..... | 1.7 | * | * | * | * | * | * | 0.7 | 2.0 | 5.4 | 19.8 | 70.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Nutritional deficiencies (E40-E64)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|-----|------|------|
| 2000..... | 1.3 | * | * | * | * | * | * | * | 2.9 | 6.2 | 17.3 | 58.5 |
| 1999..... | 1.5 | * | * | * | * | * | * | * | 3.0 | 7.8 | 22.2 | 69.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|-----|------|------|
| 2000..... | 1.8 | * | * | * | * | * | * | * | * | 3.5 | 15.9 | 70.7 |
| 1999..... | 1.9 | * | * | * | * | * | * | * | * | 3.6 | 18.4 | 71.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malnutrition (E40-E46) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.3 | 9.7 | 45.7 |
| 1999..... | 1.5 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 9.5 | 49.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 8.7 | 38.6 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 9.3 | 42.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.2 | 10.4 | 48.8 |
| 1999..... | 1.9 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 2.0 | 9.7 | 52.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 9.4 | 44.7 |
| 1999..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.9 | 8.9 | 48.8 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 8.3 | 37.9 |
| 1999..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 8.5 | 40.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.0 | 10.1 | 47.6 |
| 1999..... | 2.0 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.8 | 9.2 | 52.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | * | * | 0.6 | 1.5 | 4.5 | 15.2 | 64.0 |
| 1999..... | 1.6 | * | * | * | * | * | * | 0.6 | 1.8 | 5.2 | 18.2 | 66.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malnutrition (E40-E46) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 2.5 | 6.0 | 16.8 | 56.4 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | 2.7 | 7.5 | 19.8 | 68.5 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.6 | * | * | * | * | * | * | * | * | 3.5 | 14.3 | 67.1 |
| 1999..... | 1.7 | * | * | * | * | * | * | * | * | 3.6 | 17.3 | 65.2 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Meningitis (G00,G03) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | 2.4 | 0.2 | 0.1 | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 1.0 | 1.2 |
| 1999..... | 0.3 | 3.0 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 1.0 | 1.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | 2.4 | 0.3 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.2 | * |
| 1999..... | 0.3 | 3.1 | 0.3 | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.4 | 0.7 | 1.0 | 1.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | 2.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.0 |
| 1999..... | 0.3 | 2.9 | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.5 | 1.0 | 1.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.2 | 1.8 | 0.2 | 0.1 | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.0 |
| 1999..... | 0.3 | 2.2 | 0.3 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 0.9 | 1.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | 2.0 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.2 | * |
| 1999..... | 0.3 | 3.0 | 0.3 | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.9 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | 1.6 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.0 |
| 1999..... | 0.3 | 1.5 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.6 | 6.0 | * | * | * | * | 0.5 | 0.6 | 0.9 | 1.6 | * | * |
| 1999..... | 0.6 | 7.4 | * | * | * | * | 0.4 | 0.9 | * | 1.3 | 2.8 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Parkinson's disease (G20-G21) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.7 | * | * | * | * | * | * | 0.1 | 1.1 | 11.6 | 62.1 | 130.1 |
| 1999..... | 5.4 | * | * | * | * | * | * | 0.1 | 1.0 | 11.1 | 58.5 | 123.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.5 | * | * | * | * | * | * | 0.2 | 1.5 | 16.3 | 91.3 | 210.9 |
| 1999..... | 6.2 | * | * | * | * | * | * | 0.2 | 1.4 | 15.9 | 88.4 | 197.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.9 | * | * | * | * | * | * | 0.1 | 0.8 | 7.7 | 42.5 | 95.3 |
| 1999..... | 4.5 | * | * | * | * | * | * | * | 0.7 | 7.2 | 38.5 | 92.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.6 | * | * | * | * | * | * | 0.1 | 1.2 | 12.5 | 65.8 | 138.0 |
| 1999..... | 6.2 | * | * | * | * | * | * | 0.1 | 1.1 | 11.9 | 61.9 | 130.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 7.5 | * | * | * | * | * | * | 0.2 | 1.6 | 17.4 | 95.8 | 224.5 |
| 1999..... | 7.1 | * | * | * | * | * | * | 0.2 | 1.5 | 17.1 | 93.4 | 210.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 5.7 | * | * | * | * | * | * | * | 0.9 | 8.3 | 45.4 | 101.0 |
| 1999..... | 5.3 | * | * | * | * | * | * | * | 0.7 | 7.6 | 40.7 | 97.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | * | * | * | * | 5.5 | 27.9 | 51.7 |
| 1999..... | 1.5 | * | * | * | * | * | * | * | 0.9 | 5.8 | 24.9 | 52.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Parkinson's disease (G20-G21)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|-----|------|------|
| 2000..... | 1.8 | * | * | * | * | * | * | * | * | 8.4 | 46.5 | 79.8 |
| 1999..... | 1.6 | * | * | * | * | * | * | * | * | 8.3 | 36.9 | 80.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|-----|------|------|
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 3.5 | 16.6 | 40.0 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 3.9 | 17.6 | 40.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Alzheimer's disease (G30) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 18.0 | * | * | * | * | * | * | 0.2 | 2.0 | 18.9 | 140.1 | 659.0 |
| 1999..... | 16.3 | * | * | * | * | * | * | 0.2 | 1.9 | 17.6 | 130.4 | 598.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 10.7 | * | * | * | * | * | * | 0.2 | 1.8 | 18.9 | 125.5 | 495.4 |
| 1999..... | 10.0 | * | * | * | * | * | * | 0.2 | 2.0 | 17.5 | 121.2 | 467.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 25.0 | * | * | * | * | * | * | 0.2 | 2.2 | 18.8 | 150.0 | 729.5 |
| 1999..... | 22.3 | * | * | * | * | * | * | 0.2 | 1.8 | 17.6 | 136.5 | 653.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 20.5 | * | * | * | * | * | * | 0.2 | 2.2 | 19.8 | 145.2 | 687.6 |
| 1999..... | 18.6 | * | * | * | * | * | * | 0.2 | 2.1 | 18.3 | 135.3 | 626.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 12.2 | * | * | * | * | * | * | 0.2 | 1.9 | 19.8 | 130.3 | 517.1 |
| 1999..... | 11.4 | * | * | * | * | * | * | 0.2 | 2.1 | 18.2 | 126.0 | 491.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 28.6 | * | * | * | * | * | * | 0.2 | 2.4 | 19.8 | 155.3 | 760.5 |
| 1999..... | 25.6 | * | * | * | * | * | * | 0.2 | 2.0 | 18.5 | 141.5 | 683.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 7.7 | * | * | * | * | * | * | * | 1.2 | 15.4 | 112.0 | 446.3 |
| 1999..... | 6.7 | * | * | * | * | * | * | * | 1.2 | 14.8 | 100.1 | 373.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Alzheimer's disease (G30)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|------|------|-------|
| 2000..... | 4.6 | * | * | * | * | * | * | * | * | 15.8 | 94.7 | 342.7 |
| 1999..... | 4.2 | * | * | * | * | * | * | * | * | 15.8 | 87.6 | 294.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|---|---|---|---|------|-------|-------|
| 2000..... | 10.5 | * | * | * | * | * | * | * | * | 15.1 | 122.5 | 489.6 |
| 1999..... | 9.0 | * | * | * | * | * | * | * | * | 14.1 | 107.6 | 405.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Major cardiovascular diseases (I00-I78) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 340.4 | 16.5 | 1.5 | 0.9 | 3.4 | 10.1 | 37.0 | 117.3 | 323.2 | 859.7 | 2,405.5 | 7,925.3 |
| 1999..... | 348.5 | 16.9 | 1.5 | 0.9 | 3.5 | 10.2 | 38.0 | 118.7 | 332.8 | 897.1 | 2,490.8 | 8,155.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 323.8 | 17.4 | 1.7 | 1.0 | 4.0 | 12.8 | 49.8 | 168.0 | 449.7 | 1,127.6 | 2,877.1 | 8,015.6 |
| 1999..... | 331.5 | 17.8 | 1.6 | 1.0 | 4.1 | 13.0 | 51.5 | 170.0 | 462.1 | 1,181.9 | 2,988.4 | 8,277.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 356.2 | 15.7 | 1.4 | 0.8 | 2.8 | 7.5 | 24.4 | 68.7 | 207.8 | 639.1 | 2,087.1 | 7,886.5 |
| 1999..... | 364.7 | 15.9 | 1.4 | 0.8 | 2.8 | 7.5 | 24.6 | 69.8 | 215.0 | 663.9 | 2,157.7 | 8,104.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 360.8 | 14.3 | 1.3 | 0.8 | 2.8 | 8.4 | 31.7 | 102.9 | 295.4 | 825.9 | 2,376.9 | 8,043.5 |
| 1999..... | 369.3 | 14.3 | 1.3 | 0.8 | 3.0 | 8.5 | 32.2 | 104.4 | 304.2 | 862.2 | 2,464.4 | 8,283.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 340.9 | 15.0 | 1.6 | 0.9 | 3.4 | 11.0 | 44.2 | 151.9 | 416.9 | 1,097.1 | 2,863.9 | 8,188.3 |
| 1999..... | 348.5 | 15.1 | 1.5 | 0.9 | 3.4 | 11.1 | 45.2 | 153.5 | 428.0 | 1,149.9 | 2,980.7 | 8,447.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 379.9 | 13.6 | 1.1 | 0.6 | 2.2 | 5.8 | 19.0 | 54.9 | 182.0 | 598.9 | 2,045.9 | 7,981.6 |
| 1999..... | 389.3 | 13.5 | 1.2 | 0.7 | 2.4 | 5.9 | 19.0 | 56.5 | 188.9 | 623.0 | 2,116.6 | 8,214.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 295.4 | 29.4 | 2.8 | 1.5 | 6.8 | 22.4 | 78.1 | 247.1 | 608.0 | 1,283.0 | 3,024.3 | 7,538.2 |
| 1999..... | 300.3 | 31.1 | 2.7 | 1.5 | 6.9 | 21.5 | 82.4 | 250.0 | 624.9 | 1,340.0 | 3,083.1 | 7,597.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Major cardiovascular diseases (I00-I78)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-------|------|-----|-----|-----|------|-------|-------|-------|---------|---------|---------|
| 2000..... | 286.0 | 30.7 | 2.4 | 1.5 | 7.5 | 26.5 | 96.5 | 330.8 | 832.4 | 1,567.4 | 3,400.3 | 7,093.9 |
| 1999..... | 293.4 | 32.5 | 2.8 | 1.7 | 8.5 | 26.3 | 103.7 | 339.3 | 860.4 | 1,656.5 | 3,441.4 | 7,345.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-------|------|-----|-----|-----|------|------|-------|-------|---------|---------|---------|
| 2000..... | 303.9 | 28.0 | 3.3 | 1.6 | 6.2 | 18.7 | 61.7 | 177.3 | 436.9 | 1,075.9 | 2,795.8 | 7,723.8 |
| 1999..... | 306.6 | 29.7 | 2.6 | 1.3 | 5.2 | 17.0 | 63.7 | 175.7 | 445.3 | 1,109.4 | 2,868.2 | 7,702.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Diseases of heart (I00-I09,I11,I13,I20-I51) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 258.2 | 12.8 | 1.2 | 0.7 | 2.7 | 7.9 | 29.4 | 95.5 | 264.5 | 673.8 | 1,787.1 | 5,848.7 |
| 1999..... | 265.9 | 13.7 | 1.2 | 0.7 | 2.8 | 8.1 | 30.3 | 97.7 | 274.3 | 709.5 | 1,861.8 | 6,032.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 256.1 | 13.2 | 1.4 | 0.8 | 3.2 | 10.3 | 41.6 | 142.7 | 378.6 | 909.2 | 2,210.1 | 6,100.8 |
| 1999..... | 263.8 | 13.8 | 1.3 | 0.8 | 3.4 | 10.6 | 43.3 | 145.7 | 391.6 | 961.6 | 2,308.9 | 6,313.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 260.2 | 12.4 | 1.0 | 0.6 | 2.1 | 5.5 | 17.2 | 50.3 | 160.4 | 479.9 | 1,501.5 | 5,740.1 |
| 1999..... | 268.0 | 13.6 | 1.1 | 0.6 | 2.2 | 5.6 | 17.6 | 51.9 | 167.5 | 503.2 | 1,562.5 | 5,913.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 274.8 | 11.3 | 1.0 | 0.6 | 2.2 | 6.5 | 25.8 | 86.2 | 245.4 | 651.8 | 1,772.4 | 5,953.9 |
| 1999..... | 282.8 | 11.6 | 1.0 | 0.6 | 2.3 | 6.7 | 26.3 | 88.2 | 254.7 | 686.6 | 1,848.4 | 6,140.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 271.2 | 11.7 | 1.3 | 0.7 | 2.6 | 8.8 | 37.9 | 132.2 | 356.2 | 891.2 | 2,209.6 | 6,257.6 |
| 1999..... | 278.8 | 11.8 | 1.2 | 0.7 | 2.8 | 9.0 | 38.8 | 134.7 | 367.6 | 942.1 | 2,313.9 | 6,462.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 278.3 | 10.9 | 0.8 | 0.4 | 1.6 | 4.2 | 13.7 | 41.2 | 142.2 | 451.3 | 1,475.2 | 5,824.0 |
| 1999..... | 286.6 | 11.4 | 0.9 | 0.6 | 1.9 | 4.4 | 13.8 | 42.8 | 149.5 | 474.3 | 1,534.8 | 6,006.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 219.6 | 20.9 | 2.2 | 1.2 | 5.7 | 17.9 | 57.9 | 186.0 | 472.0 | 977.3 | 2,200.0 | 5,403.4 |
| 1999..... | 225.4 | 25.3 | 2.2 | 1.2 | 5.9 | 17.3 | 62.7 | 191.5 | 489.3 | 1,027.1 | 2,253.8 | 5,498.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Diseases of heart (I00-I09,I11,I13,I20-I51)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-------|------|-----|-----|-----|------|------|-------|-------|---------|---------|---------|
| 2000..... | 219.0 | 20.9 | 2.1 | 1.3 | 6.6 | 22.0 | 74.9 | 257.5 | 659.4 | 1,212.8 | 2,522.4 | 5,198.6 |
| 1999..... | 226.9 | 24.9 | 2.4 | 1.4 | 7.5 | 21.8 | 82.6 | 267.6 | 690.3 | 1,294.0 | 2,560.2 | 5,433.3 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-------|------|-----|-----|-----|------|------|-------|-------|-------|---------|---------|
| 2000..... | 220.1 | 21.0 | 2.3 | 1.1 | 4.8 | 14.2 | 42.9 | 126.3 | 329.1 | 805.9 | 2,004.2 | 5,489.0 |
| 1999..... | 224.0 | 25.7 | 2.1 | 1.0 | 4.4 | 13.2 | 45.1 | 128.3 | 336.1 | 832.7 | 2,070.1 | 5,525.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 4.5 | 9.6 | 16.6 |
| 1999..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 4.7 | 10.1 | 17.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.8 | * | * | * | * | * | 0.3 | 0.6 | 1.5 | 2.9 | 6.5 | 11.6 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 6.7 | 9.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.8 | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.0 | 5.8 | 11.6 | 18.7 |
| 1999..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 5.6 | 12.4 | 21.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.7 | 4.6 | 10.2 | 17.4 |
| 1999..... | 1.5 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 4.9 | 10.6 | 18.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 2.9 | 6.9 | 12.1 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.7 | 6.9 | 10.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | * | * | * | 0.3 | 0.7 | 1.9 | 6.1 | 12.4 | 19.7 |
| 1999..... | 2.1 | * | * | * | * | * | 0.3 | 0.8 | 2.2 | 5.8 | 13.1 | 22.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.8 | * | * | * | * | * | 0.6 | 1.0 | 2.1 | 3.4 | 4.0 | 8.5 |
| 1999..... | 0.8 | * | * | * | * | * | 0.6 | 1.2 | 2.2 | 3.3 | 4.5 | 8.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

BLACK, MALE

| | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|---|---|
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|------|
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | 1.3 | 2.6 | 4.3 | 5.0 | * |
| 1999..... | 1.0 | * | * | * | * | * | 0.7 | 1.4 | 2.6 | 4.1 | 4.7 | 11.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Hypertensive heart disease (I11) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 8.6 | * | * | * | 0.1 | 0.6 | 2.2 | 5.7 | 11.1 | 19.8 | 48.4 | 190.6 |
| 1999..... | 8.3 | * | * | * | 0.1 | 0.6 | 2.2 | 5.4 | 10.8 | 19.3 | 48.0 | 183.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 7.3 | * | * | * | 0.1 | 0.7 | 2.9 | 7.9 | 14.7 | 24.0 | 45.7 | 132.0 |
| 1999..... | 7.0 | * | * | * | 0.1 | 0.8 | 3.0 | 7.2 | 14.0 | 22.1 | 46.2 | 128.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 9.9 | * | * | * | * | 0.4 | 1.5 | 3.5 | 7.9 | 16.4 | 50.2 | 215.8 |
| 1999..... | 9.6 | * | * | * | * | 0.3 | 1.3 | 3.7 | 8.0 | 17.0 | 49.2 | 207.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 7.7 | * | * | * | * | 0.3 | 1.5 | 3.7 | 7.7 | 15.2 | 41.7 | 179.8 |
| 1999..... | 7.4 | * | * | * | * | 0.4 | 1.3 | 3.5 | 7.6 | 14.6 | 41.2 | 175.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 6.2 | * | * | * | * | 0.5 | 2.1 | 5.6 | 10.5 | 18.9 | 38.9 | 123.4 |
| 1999..... | 5.9 | * | * | * | * | 0.6 | 2.0 | 5.0 | 10.2 | 16.9 | 39.3 | 119.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 9.1 | * | * | * | * | 0.2 | 0.9 | 1.9 | 5.1 | 12.1 | 43.5 | 203.9 |
| 1999..... | 8.9 | * | * | * | * | * | 0.7 | 2.0 | 5.2 | 12.7 | 42.5 | 198.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.7 | * | * | * | * | 2.2 | 7.4 | 21.7 | 42.7 | 65.2 | 134.3 | 340.9 |
| 1999..... | 16.1 | * | * | * | 0.3 | 1.9 | 8.0 | 21.6 | 40.3 | 65.8 | 134.4 | 305.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Hypertensive heart disease (I11)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|------|------|------|------|-------|-------|
| 2000..... | 16.2 | * | * | * | * | 2.7 | 9.2 | 28.9 | 57.4 | 77.8 | 135.2 | 252.2 |
| 1999..... | 15.7 | * | * | * | * | 2.2 | 10.2 | 27.6 | 52.9 | 77.3 | 138.9 | 244.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|------|------|------|-------|-------|
| 2000..... | 17.1 | * | * | * | * | 1.7 | 5.8 | 15.6 | 31.6 | 56.0 | 133.8 | 378.0 |
| 1999..... | 16.5 | * | * | * | * | 1.5 | 6.0 | 16.6 | 30.7 | 57.5 | 131.7 | 331.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Hypertensive heart and renal disease (I13) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 7.0 | 24.0 |
| 1999..... | 1.2 | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.1 | 2.6 | 8.9 | 29.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.5 | 7.1 | 26.4 |
| 1999..... | 1.0 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.7 | 9.4 | 30.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.1 | 7.0 | 23.0 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.5 | 8.6 | 28.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.5 | 5.6 | 21.8 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 7.4 | 26.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.7 | 6.1 | 23.2 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 8.1 | 27.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 5.3 | 21.2 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 6.8 | 25.2 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.3 | * | * | * | * | * | 0.8 | 2.2 | 4.7 | 9.7 | 23.2 | 50.5 |
| 1999..... | 2.9 | * | * | * | * | * | 1.5 | 2.5 | 6.1 | 12.2 | 27.4 | 65.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Hypertensive heart and renal disease (I13) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 2.1 | * | * | * | * | * | 1.0 | 2.9 | 5.6 | 10.8 | 20.3 | 58.5 |
| 1999..... | 2.6 | * | * | * | * | * | 1.8 | 3.6 | 8.0 | 13.0 | 25.2 | 55.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.4 | * | * | * | * | * | * | 1.6 | 4.0 | 8.9 | 25.0 | 47.1 |
| 1999..... | 3.1 | * | * | * | * | * | 1.2 | 1.5 | 4.7 | 11.6 | 28.8 | 69.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 187.2 | 0.8 | * | * | 0.3 | 2.5 | 16.7 | 67.0 | 199.1 | 511.3 | 1,329.4 | 4,127.8 |
| 1999..... | 194.2 | 0.7 | * | 0.1 | 0.4 | 2.7 | 17.3 | 69.2 | 206.8 | 542.5 | 1,393.8 | 4,296.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 193.6 | * | * | * | 0.4 | 3.8 | 25.9 | 105.4 | 295.6 | 713.7 | 1,707.5 | 4,461.0 |
| 1999..... | 200.5 | * | * | * | 0.5 | 3.9 | 26.8 | 108.9 | 306.0 | 758.5 | 1,792.3 | 4,654.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 181.1 | * | * | * | 0.2 | 1.2 | 7.7 | 30.2 | 111.2 | 344.6 | 1,074.0 | 3,984.3 |
| 1999..... | 188.2 | * | * | * | 0.2 | 1.4 | 7.9 | 31.2 | 116.5 | 365.8 | 1,126.9 | 4,145.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 201.3 | 0.8 | * | * | 0.3 | 2.3 | 15.9 | 63.4 | 189.0 | 500.5 | 1,324.3 | 4,210.4 |
| 1999..... | 208.9 | * | * | * | 0.3 | 2.5 | 16.2 | 65.5 | 197.0 | 531.7 | 1,391.9 | 4,380.8 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 207.7 | * | * | * | 0.4 | 3.5 | 25.2 | 101.7 | 283.8 | 707.8 | 1,715.7 | 4,591.4 |
| 1999..... | 215.0 | * | * | * | 0.4 | 3.6 | 25.6 | 104.8 | 294.0 | 752.3 | 1,808.3 | 4,780.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 195.1 | * | * | * | 0.2 | 1.0 | 6.5 | 25.8 | 100.7 | 326.9 | 1,058.3 | 4,047.5 |
| 1999..... | 203.0 | * | * | * | 0.2 | 1.3 | 6.8 | 27.1 | 106.7 | 348.4 | 1,111.3 | 4,213.2 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 145.2 | * | * | * | 0.4 | 4.4 | 25.8 | 108.2 | 318.6 | 684.5 | 1,551.6 | 3,694.1 |
| 1999..... | 149.1 | * | * | * | 0.8 | 4.0 | 27.8 | 112.1 | 325.4 | 720.5 | 1,582.6 | 3,813.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 146.8 | * | * | * | * | 6.1 | 35.9 | 155.7 | 458.8 | 867.8 | 1,822.6 | 3,585.3 |
| 1999..... | 151.6 | * | * | * | 0.9 | 5.9 | 39.8 | 164.7 | 473.7 | 924.8 | 1,825.7 | 3,783.8 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 143.8 | * | * | * | * | 2.9 | 16.9 | 68.6 | 211.7 | 551.1 | 1,387.0 | 3,739.6 |
| 1999..... | 146.9 | * | * | * | 0.7 | 2.3 | 17.1 | 68.3 | 212.4 | 571.7 | 1,436.9 | 3,825.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Acute myocardial infarction (I21-I22) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 70.1 | 0.5 | * | * | 0.1 | 1.0 | 7.0 | 29.3 | 87.7 | 211.7 | 505.2 | 1,321.0 |
| 1999..... | 73.1 | * | * | * | 0.2 | 1.1 | 7.3 | 30.2 | 91.5 | 227.5 | 532.7 | 1,372.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 74.5 | * | * | * | 0.2 | 1.4 | 10.7 | 45.5 | 128.0 | 290.4 | 637.3 | 1,488.0 |
| 1999..... | 78.2 | * | * | * | 0.2 | 1.6 | 11.1 | 47.4 | 134.8 | 313.6 | 676.3 | 1,560.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 65.8 | * | * | * | 0.1 | 0.6 | 3.3 | 13.8 | 50.8 | 146.9 | 416.0 | 1,249.0 |
| 1999..... | 68.3 | * | * | * | 0.1 | 0.7 | 3.5 | 13.8 | 52.1 | 157.0 | 436.6 | 1,292.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 75.4 | * | * | * | 0.1 | 0.9 | 6.8 | 28.4 | 85.7 | 209.7 | 503.9 | 1,338.4 |
| 1999..... | 78.8 | * | * | * | 0.1 | 1.1 | 7.0 | 29.6 | 89.8 | 226.2 | 533.3 | 1,390.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 80.4 | * | * | * | 0.2 | 1.4 | 10.6 | 45.2 | 126.6 | 291.5 | 641.6 | 1,519.3 |
| 1999..... | 84.5 | * | * | * | 0.2 | 1.5 | 11.0 | 47.3 | 133.5 | 315.7 | 684.9 | 1,593.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 70.6 | * | * | * | * | 0.5 | 2.9 | 12.0 | 47.6 | 141.2 | 410.4 | 1,261.1 |
| 1999..... | 73.4 | * | * | * | * | 0.6 | 3.1 | 12.3 | 49.1 | 151.9 | 431.2 | 1,305.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 54.1 | * | * | * | * | 1.8 | 10.0 | 42.1 | 121.7 | 262.0 | 581.7 | 1,278.9 |
| 1999..... | 55.3 | * | * | * | * | 1.7 | 10.4 | 42.2 | 122.9 | 273.8 | 595.9 | 1,321.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Acute myocardial infarction (I21-I22)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|------|------|-------|-------|-------|---------|
| 2000..... | 53.9 | * | * | * | * | 2.3 | 13.5 | 57.2 | 167.5 | 321.0 | 667.4 | 1,309.0 |
| 1999..... | 55.5 | * | * | * | * | 2.4 | 14.3 | 59.4 | 173.1 | 339.8 | 669.1 | 1,389.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|------|------|-------|-------|---------|
| 2000..... | 54.3 | * | * | * | * | 1.4 | 6.8 | 29.4 | 86.7 | 219.0 | 529.7 | 1,266.4 |
| 1999..... | 55.1 | * | * | * | * | 1.0 | 6.9 | 27.9 | 84.6 | 225.8 | 552.0 | 1,292.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other acute ischemic heart diseases (I24) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | * | * | * | 0.2 | 0.8 | 2.0 | 3.5 | 7.4 | 21.6 |
| 1999..... | 1.2 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 3.7 | 7.0 | 21.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | * | * | * | 0.3 | 1.2 | 3.0 | 4.9 | 9.0 | 19.6 |
| 1999..... | 1.2 | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 5.0 | 8.7 | 21.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 2.5 | 6.3 | 22.4 |
| 1999..... | 1.1 | * | * | * | * | * | * | 0.4 | 1.1 | 2.6 | 5.9 | 21.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 3.3 | 7.1 | 21.9 |
| 1999..... | 1.2 | * | * | * | * | * | 0.1 | 0.6 | 1.8 | 3.5 | 6.8 | 22.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | * | * | * | 0.3 | 1.1 | 2.7 | 4.5 | 8.7 | 20.0 |
| 1999..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 4.7 | 8.5 | 22.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | * | * | * | * | 0.3 | 0.9 | 2.4 | 6.0 | 22.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.3 | 1.1 | 2.5 | 5.7 | 22.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | * | * | * | 0.4 | 1.5 | 4.3 | 5.5 | 11.0 | 22.0 |
| 1999..... | 1.2 | * | * | * | * | * | 0.5 | 1.9 | 3.6 | 5.5 | 10.9 | 17.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other acute ischemic heart diseases (I24)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|------|---|
| 2000..... | 1.5 | * | * | * | * | * | * | 2.4 | 6.4 | 8.3 | 13.5 | * |
| 1999..... | 1.5 | * | * | * | * | * | 0.8 | 2.7 | 6.4 | 7.9 | 12.3 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|------|------|
| 2000..... | 1.1 | * | * | * | * | * | * | * | 2.6 | 3.5 | 9.5 | 23.1 |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.1 | 1.5 | 3.8 | 10.1 | 16.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other forms of chronic ischemic heart disease (I20,I25) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 115.9 | * | * | * | 0.2 | 1.5 | 9.6 | 37.0 | 109.5 | 296.0 | 816.8 | 2,785.2 |
| 1999..... | 119.9 | * | * | * | 0.2 | 1.5 | 9.9 | 38.3 | 113.4 | 311.4 | 854.0 | 2,902.2 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 117.7 | * | * | * | 0.2 | 2.3 | 14.9 | 58.7 | 164.6 | 418.5 | 1,061.2 | 2,953.4 |
| 1999..... | 121.1 | * | * | * | 0.3 | 2.3 | 15.4 | 60.5 | 168.4 | 439.9 | 1,107.3 | 3,072.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 114.1 | * | * | * | * | 0.6 | 4.3 | 16.1 | 59.2 | 195.2 | 651.8 | 2,712.8 |
| 1999..... | 118.7 | * | * | * | 0.1 | 0.8 | 4.4 | 17.0 | 63.3 | 206.2 | 684.4 | 2,830.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 124.6 | * | * | * | 0.2 | 1.3 | 8.9 | 34.2 | 101.6 | 287.5 | 813.3 | 2,850.1 |
| 1999..... | 128.8 | * | * | * | 0.1 | 1.4 | 9.0 | 35.3 | 105.4 | 301.9 | 851.7 | 2,967.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 126.0 | * | * | * | 0.2 | 2.1 | 14.3 | 55.4 | 154.6 | 411.8 | 1,065.4 | 3,052.1 |
| 1999..... | 129.3 | * | * | * | 0.2 | 2.1 | 14.4 | 56.6 | 157.9 | 431.8 | 1,114.9 | 3,165.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 123.3 | * | * | * | * | 0.5 | 3.5 | 13.5 | 52.1 | 183.4 | 641.9 | 2,763.7 |
| 1999..... | 128.4 | * | * | * | * | 0.7 | 3.6 | 14.5 | 56.5 | 194.0 | 674.4 | 2,885.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 89.8 | * | * | * | * | 2.6 | 15.5 | 64.6 | 192.7 | 417.0 | 958.9 | 2,393.2 |
| 1999..... | 92.6 | * | * | * | 0.5 | 2.3 | 16.9 | 68.0 | 198.9 | 441.2 | 975.8 | 2,474.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other forms of chronic ischemic heart disease (I20,I25)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|------|-------|-------|-------|---------|---------|
| 2000..... | 91.3 | * | * | * | * | 3.7 | 21.8 | 96.0 | 284.9 | 538.6 | 1,141.7 | 2,257.2 |
| 1999..... | 94.6 | * | * | * | * | 3.5 | 24.8 | 102.5 | 294.2 | 577.1 | 1,144.3 | 2,375.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|------|------|-------|-------|-------|---------|
| 2000..... | 88.4 | * | * | * | * | 1.5 | 9.8 | 38.5 | 122.3 | 328.6 | 847.9 | 2,450.1 |
| 1999..... | 90.8 | * | * | * | * | 1.3 | 10.0 | 39.3 | 126.3 | 342.1 | 874.8 | 2,516.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Atherosclerotic cardiovascular disease, so described (I25.0) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 25.3 | * | * | * | * | 0.4 | 3.1 | 13.4 | 34.4 | 69.2 | 158.7 | 527.6 |
| 1999..... | 26.4 | * | * | * | * | 0.4 | 3.3 | 13.2 | 35.8 | 73.5 | 167.3 | 560.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 26.7 | * | * | * | * | 0.6 | 4.9 | 21.9 | 53.6 | 98.4 | 195.9 | 522.6 |
| 1999..... | 27.5 | * | * | * | * | 0.6 | 5.4 | 21.4 | 54.9 | 104.2 | 204.5 | 556.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 23.9 | * | * | * | * | 0.2 | 1.2 | 5.3 | 16.9 | 45.1 | 133.6 | 529.8 |
| 1999..... | 25.3 | * | * | * | * | 0.2 | 1.3 | 5.4 | 18.5 | 48.4 | 142.4 | 562.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 25.5 | * | * | * | * | 0.3 | 2.8 | 11.9 | 29.7 | 62.1 | 149.6 | 523.6 |
| 1999..... | 26.6 | * | * | * | * | 0.4 | 3.0 | 11.5 | 30.8 | 65.6 | 157.9 | 556.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 26.5 | * | * | * | * | 0.6 | 4.6 | 19.8 | 46.9 | 89.1 | 184.4 | 520.4 |
| 1999..... | 27.2 | * | * | * | * | 0.6 | 4.9 | 19.0 | 47.7 | 93.9 | 193.2 | 553.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 24.6 | * | * | * | * | * | 1.0 | 4.2 | 13.6 | 39.5 | 125.9 | 525.0 |
| 1999..... | 26.0 | * | * | * | * | 0.2 | 1.0 | 4.2 | 15.1 | 42.1 | 134.2 | 557.6 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 29.5 | * | * | * | * | 0.7 | 5.2 | 27.6 | 79.2 | 141.9 | 283.6 | 648.3 |
| 1999..... | 31.0 | * | * | * | * | 0.8 | 6.4 | 28.4 | 82.8 | 155.4 | 296.8 | 670.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Atherosclerotic cardiovascular disease, so described (I25.0)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|------|-------|-------|-------|-------|
| 2000..... | 33.5 | * | * | * | * | 1.0 | 7.8 | 43.7 | 125.3 | 198.2 | 359.0 | 647.0 |
| 1999..... | 35.1 | * | * | * | * | 1.1 | 9.7 | 45.0 | 129.0 | 220.3 | 370.8 | 664.0 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|---|-----|------|------|-------|-------|-------|
| 2000..... | 25.9 | * | * | * | * | * | 2.8 | 14.2 | 44.1 | 100.9 | 237.9 | 648.8 |
| 1999..... | 27.4 | * | * | * | * | * | 3.5 | 14.7 | 47.6 | 108.0 | 252.4 | 672.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

ALL AGES 1/
UNDER 1 YEAR 2/
1-4 YEARS
5-14 YEARS
15-24 YEARS
25-34 YEARS
35-44 YEARS
45-54 YEARS
55-64 YEARS
65-74 YEARS
75-84 YEARS
85 YEARS AND OVER

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

| | | | | | | | | | | | | |
|--------------------------|-------|---|---|---|-----|-----|------|------|-------|-------|-------|---------|
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 90.6 | * | * | * | 0.1 | 1.1 | 6.5 | 23.6 | 75.1 | 226.9 | 658.1 | 2,257.6 |
| 1999..... | 93.5 | * | * | * | 0.2 | 1.1 | 6.5 | 25.0 | 77.6 | 237.8 | 686.7 | 2,341.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 91.1 | * | * | * | 0.2 | 1.7 | 10.0 | 36.8 | 111.0 | 320.1 | 865.3 | 2,430.8 |
| 1999..... | 93.6 | * | * | * | 0.2 | 1.6 | 10.1 | 39.0 | 113.5 | 335.7 | 902.8 | 2,515.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 90.1 | * | * | * | * | 0.5 | 3.1 | 10.9 | 42.3 | 150.1 | 518.2 | 2,183.1 |
| 1999..... | 93.4 | * | * | * | 0.1 | 0.5 | 3.0 | 11.6 | 44.8 | 157.8 | 542.0 | 2,268.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 99.1 | * | * | * | 0.1 | 1.0 | 6.1 | 22.3 | 71.9 | 225.4 | 663.7 | 2,326.5 |
| 1999..... | 102.2 | * | * | * | 0.1 | 1.0 | 6.1 | 23.8 | 74.6 | 236.3 | 693.8 | 2,411.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 99.5 | * | * | * | 0.1 | 1.5 | 9.7 | 35.6 | 107.7 | 322.7 | 881.0 | 2,531.7 |
| 1999..... | 102.1 | * | * | * | 0.2 | 1.6 | 9.5 | 37.6 | 110.2 | 337.9 | 921.8 | 2,612.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 98.7 | * | * | * | * | 0.4 | 2.5 | 9.3 | 38.5 | 143.9 | 516.0 | 2,238.8 |
| 1999..... | 102.4 | * | * | * | * | 0.5 | 2.6 | 10.2 | 41.4 | 151.8 | 540.2 | 2,327.5 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 60.3 | * | * | * | * | 1.9 | 10.3 | 37.0 | 113.4 | 275.2 | 675.3 | 1,745.0 |
| 1999..... | 61.6 | * | * | * | 0.3 | 1.6 | 10.5 | 39.6 | 116.1 | 285.8 | 679.0 | 1,804.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|------|------|-------|-------|-------|---------|
| 2000..... | 57.9 | * | * | * | * | 2.7 | 14.0 | 52.3 | 159.5 | 340.4 | 782.7 | 1,610.1 |
| 1999..... | 59.5 | * | * | * | * | 2.4 | 15.1 | 57.6 | 165.2 | 356.7 | 773.5 | 1,711.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|------|------|-------|-------|---------|
| 2000..... | 62.5 | * | * | * | * | 1.2 | 7.0 | 24.3 | 78.3 | 227.7 | 610.0 | 1,801.3 |
| 1999..... | 63.4 | * | * | * | * | 0.8 | 6.5 | 24.6 | 78.7 | 234.1 | 622.4 | 1,843.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other heart diseases (I26-I51) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 60.1 | 11.9 | 1.1 | 0.6 | 2.2 | 4.7 | 9.9 | 21.8 | 51.6 | 135.8 | 392.8 | 1,489.7 |
| 1999..... | 60.8 | 12.8 | 1.2 | 0.6 | 2.3 | 4.7 | 10.4 | 22.0 | 53.8 | 140.4 | 401.0 | 1,505.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 53.6 | 12.1 | 1.3 | 0.7 | 2.6 | 5.6 | 12.3 | 28.3 | 65.9 | 166.0 | 443.4 | 1,469.8 |
| 1999..... | 54.5 | 13.1 | 1.3 | 0.7 | 2.8 | 5.7 | 13.0 | 28.5 | 69.1 | 174.7 | 454.3 | 1,490.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 66.3 | 11.6 | 0.9 | 0.5 | 1.8 | 3.7 | 7.6 | 15.5 | 38.5 | 111.0 | 358.6 | 1,498.2 |
| 1999..... | 66.9 | 12.6 | 1.0 | 0.6 | 1.9 | 3.7 | 7.8 | 15.9 | 39.9 | 112.3 | 365.4 | 1,511.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 63.5 | 10.4 | 0.9 | 0.5 | 1.8 | 3.8 | 8.1 | 18.3 | 46.6 | 129.9 | 390.6 | 1,524.5 |
| 1999..... | 64.0 | 10.9 | 1.0 | 0.6 | 2.0 | 3.8 | 8.5 | 18.5 | 47.8 | 133.9 | 397.4 | 1,540.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 55.7 | 10.6 | 1.2 | 0.6 | 2.1 | 4.7 | 10.3 | 24.2 | 59.9 | 159.8 | 442.1 | 1,507.5 |
| 1999..... | 56.2 | 11.2 | 1.1 | 0.6 | 2.3 | 4.7 | 10.9 | 24.2 | 61.7 | 167.6 | 451.2 | 1,522.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 71.1 | 10.2 | 0.7 | 0.4 | 1.4 | 2.9 | 6.0 | 12.6 | 34.1 | 104.8 | 355.6 | 1,531.7 |
| 1999..... | 71.5 | 10.5 | 0.8 | 0.5 | 1.7 | 2.9 | 6.0 | 12.8 | 34.9 | 106.0 | 361.2 | 1,547.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 54.7 | 20.1 | 2.2 | 1.2 | 4.8 | 10.8 | 23.3 | 52.8 | 103.9 | 214.6 | 486.8 | 1,309.4 |
| 1999..... | 56.5 | 23.7 | 2.2 | 1.1 | 4.6 | 10.9 | 24.9 | 54.2 | 115.2 | 225.2 | 504.8 | 1,305.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other heart diseases (I26-I51) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 53.4 | 19.9 | 2.1 | 1.3 | 5.4 | 12.7 | 28.3 | 69.3 | 136.3 | 254.2 | 541.9 | 1,294.1 |
| 1999..... | 56.4 | 23.2 | 2.4 | 1.2 | 5.8 | 13.2 | 30.3 | 70.6 | 154.0 | 276.6 | 566.2 | 1,348.5 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 55.8 | 20.3 | 2.2 | 1.1 | 4.2 | 9.0 | 18.8 | 39.1 | 79.1 | 185.8 | 453.4 | 1,315.8 |
| 1999..... | 56.5 | 24.3 | 2.0 | 0.9 | 3.4 | 8.8 | 20.1 | 40.5 | 85.6 | 187.8 | 468.0 | 1,287.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Acute and subacute endocarditis (I33) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.2 | 2.5 | 2.8 |
| 1999..... | 0.4 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.3 | 2.4 | 2.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 0.8 | 1.5 | 3.4 | 3.9 |
| 1999..... | 0.5 | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 1.4 | 3.2 | 4.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 1.8 | 2.3 |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 1.9 | 2.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.2 | 2.4 | 2.7 |
| 1999..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.4 | 3.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.4 | 3.4 | 4.0 |
| 1999..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.4 | 3.3 | 4.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.0 | 1.8 | 2.1 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 1.8 | 2.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.6 | * | * | * | * | * | 0.5 | 1.4 | 1.6 | 1.8 | 2.9 | * |
| 1999..... | 0.6 | * | * | * | * | * | 0.7 | 1.2 | 2.0 | 2.5 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Acute and subacute endocarditis (I33)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|---|---|---|
| 2000..... | 0.7 | * | * | * | * | * | 0.8 | 1.9 | * | * | * | * |
| 1999..... | 0.7 | * | * | * | * | * | 1.0 | 2.0 | 2.7 | * | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|---|---|
| 2000..... | 0.6 | * | * | * | * | * | * | 1.0 | 1.5 | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 1.6 | 2.8 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Diseases of pericardium and acute myocarditis (I30-I31,I40) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | 0.5 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.7 | 1.2 | 1.8 |
| 1999..... | 0.3 | 0.7 | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.3 | 0.5 | 0.8 | 1.3 | 2.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.5 | 2.0 |
| 1999..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.6 | 0.8 | 1.3 | 2.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.0 | 1.7 |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 1.3 | 1.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.2 | 1.8 |
| 1999..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.3 | 2.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.7 | 1.4 | 2.0 |
| 1999..... | 0.3 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 0.8 | 1.4 | 2.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 1.7 |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.3 | 1.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | * | 0.4 | 0.5 | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | 0.5 | 0.5 | * | 1.2 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Heart failure (I50) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 20.2 | 0.8 | * | 0.1 | 0.1 | 0.2 | 0.6 | 2.2 | 9.2 | 35.2 | 136.6 | 675.9 |
| 1999..... | 20.1 | * | * | * | 0.1 | 0.2 | 0.6 | 2.2 | 9.0 | 35.8 | 138.1 | 678.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 15.7 | 1.1 | * | * | 0.1 | 0.2 | 0.7 | 2.7 | 11.1 | 41.6 | 151.1 | 637.8 |
| 1999..... | 15.6 | * | * | * | 0.1 | 0.2 | 0.7 | 2.8 | 11.0 | 43.0 | 150.2 | 641.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 24.6 | * | * | * | * | 0.1 | 0.5 | 1.7 | 7.6 | 30.0 | 126.8 | 692.3 |
| 1999..... | 24.5 | * | * | * | * | 0.2 | 0.5 | 1.6 | 7.2 | 29.9 | 129.9 | 694.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 22.2 | * | * | * | * | 0.1 | 0.4 | 1.8 | 8.3 | 33.9 | 137.0 | 695.9 |
| 1999..... | 22.0 | * | * | * | * | 0.1 | 0.5 | 1.7 | 8.1 | 34.5 | 137.5 | 697.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 17.0 | * | * | * | * | 0.1 | 0.6 | 2.3 | 10.1 | 40.7 | 152.0 | 658.2 |
| 1999..... | 16.7 | * | * | * | * | * | 0.6 | 2.2 | 10.0 | 41.9 | 150.5 | 658.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 27.1 | * | * | * | * | * | 0.3 | 1.3 | 6.6 | 28.3 | 126.8 | 712.1 |
| 1999..... | 27.0 | * | * | * | * | * | 0.3 | 1.2 | 6.3 | 28.4 | 128.8 | 714.2 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 14.1 | * | * | * | * | 0.5 | 1.6 | 5.8 | 18.9 | 54.0 | 157.3 | 559.8 |
| 1999..... | 14.3 | * | * | * | * | 0.6 | 1.7 | 6.6 | 19.2 | 55.2 | 166.6 | 554.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Heart failure (I50) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 11.6 | * | * | * | * | * | 1.8 | 7.3 | 22.3 | 59.2 | 168.5 | 536.4 |
| 1999..... | 12.2 | * | * | * | * | * | 1.9 | 8.8 | 23.9 | 62.6 | 174.9 | 559.6 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 16.3 | * | * | * | * | * | 1.6 | 4.6 | 16.2 | 50.3 | 150.5 | 569.6 |
| 1999..... | 16.3 | * | * | * | * | * | 1.4 | 4.8 | 15.6 | 49.8 | 161.6 | 551.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 39.2 | 10.5 | 0.9 | 0.5 | 2.0 | 4.3 | 8.9 | 18.9 | 41.3 | 98.6 | 252.6 | 809.2 |
| 1999..... | 39.9 | 11.8 | 0.9 | 0.5 | 2.1 | 4.3 | 9.3 | 19.2 | 43.5 | 102.4 | 259.3 | 821.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 37.0 | 10.4 | 1.0 | 0.6 | 2.4 | 5.2 | 11.0 | 24.7 | 53.5 | 122.1 | 287.4 | 826.1 |
| 1999..... | 38.1 | 12.0 | 1.0 | 0.6 | 2.5 | 5.2 | 11.6 | 24.8 | 56.5 | 129.4 | 299.5 | 841.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 41.2 | 10.6 | 0.8 | 0.4 | 1.6 | 3.4 | 6.8 | 13.3 | 30.1 | 79.3 | 229.0 | 802.0 |
| 1999..... | 41.7 | 11.5 | 0.9 | 0.5 | 1.8 | 3.4 | 6.9 | 13.8 | 31.7 | 80.4 | 232.3 | 813.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 40.7 | 9.4 | 0.8 | 0.4 | 1.6 | 3.5 | 7.3 | 16.0 | 37.3 | 94.1 | 250.0 | 824.1 |
| 1999..... | 41.4 | 10.2 | 0.8 | 0.5 | 1.8 | 3.5 | 7.6 | 16.2 | 38.6 | 97.4 | 256.2 | 837.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 37.9 | 9.2 | 0.9 | 0.6 | 2.0 | 4.3 | 9.2 | 21.2 | 48.6 | 117.0 | 285.2 | 843.3 |
| 1999..... | 38.7 | 10.5 | 1.0 | 0.5 | 2.1 | 4.3 | 9.8 | 21.3 | 50.3 | 123.4 | 296.0 | 856.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 43.4 | 9.5 | 0.6 | 0.3 | 1.2 | 2.6 | 5.4 | 10.9 | 26.8 | 74.9 | 226.1 | 815.9 |
| 1999..... | 43.9 | 9.8 | 0.7 | 0.5 | 1.6 | 2.7 | 5.4 | 11.2 | 27.7 | 75.8 | 229.3 | 829.5 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 39.6 | 16.5 | 1.6 | 1.0 | 4.3 | 9.8 | 20.6 | 45.1 | 82.8 | 157.7 | 325.5 | 743.9 |
| 1999..... | 41.1 | 21.4 | 1.6 | 0.9 | 4.3 | 10.0 | 22.0 | 45.8 | 93.2 | 166.3 | 334.5 | 747.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|------|------|------|-------|-------|-------|-------|
| 2000..... | 40.7 | 15.5 | 1.8 | 1.1 | 4.9 | 11.6 | 25.2 | 59.6 | 111.7 | 191.3 | 368.7 | 753.4 |
| 1999..... | 43.1 | 21.1 | * | 1.1 | 5.4 | 12.0 | 26.9 | 59.2 | 126.5 | 210.3 | 388.5 | 787.8 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|---|-----|-----|-----|------|------|------|-------|-------|-------|
| 2000..... | 38.6 | 17.5 | * | 0.8 | 3.8 | 8.1 | 16.6 | 33.0 | 60.7 | 133.2 | 299.3 | 739.9 |
| 1999..... | 39.3 | 21.8 | * | * | 3.1 | 8.2 | 17.7 | 34.6 | 67.8 | 134.2 | 302.1 | 730.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Essential (primary) hypertension and hypertensive renal disease (I10,I12) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.6 | * | * | * | * | 0.2 | 0.8 | 2.4 | 6.0 | 15.3 | 45.7 | 160.8 |
| 1999..... | 6.2 | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.6 | 15.4 | 43.8 | 151.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 5.1 | * | * | * | * | 0.2 | 1.0 | 3.1 | 7.2 | 15.9 | 44.4 | 135.1 |
| 1999..... | 4.9 | * | * | * | * | 0.3 | 0.9 | 2.8 | 6.6 | 16.7 | 42.4 | 127.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 7.9 | * | * | * | * | 0.2 | 0.6 | 1.7 | 5.0 | 14.8 | 46.5 | 171.9 |
| 1999..... | 7.5 | * | * | * | * | 0.2 | 0.5 | 1.7 | 4.8 | 14.2 | 44.8 | 161.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.1 | * | * | * | * | 0.1 | 0.4 | 1.4 | 4.1 | 12.0 | 40.5 | 151.5 |
| 1999..... | 5.7 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.8 | 12.2 | 38.2 | 141.8 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 4.6 | * | * | * | * | 0.1 | 0.6 | 1.9 | 5.1 | 12.8 | 39.7 | 126.4 |
| 1999..... | 4.3 | * | * | * | * | 0.2 | 0.5 | 1.7 | 4.6 | 13.6 | 36.9 | 117.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 7.5 | * | * | * | * | * | 0.3 | 0.9 | 3.3 | 11.3 | 41.1 | 162.3 |
| 1999..... | 7.1 | * | * | * | * | * | 0.2 | 1.0 | 3.0 | 11.1 | 39.0 | 152.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.0 | * | * | * | * | 0.7 | 3.1 | 9.9 | 23.2 | 47.5 | 109.1 | 284.1 |
| 1999..... | 10.7 | * | * | * | * | 0.7 | 3.1 | 9.4 | 21.9 | 45.7 | 113.8 | 270.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|------|------|-------|-------|
| 2000..... | 9.5 | * | * | * | * | * | 3.7 | 12.4 | 29.0 | 48.2 | 105.8 | 249.0 |
| 1999..... | 9.5 | * | * | * | * | * | 3.7 | 12.2 | 25.7 | 49.3 | 115.8 | 246.7 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|-----|-----|------|------|-------|-------|
| 2000..... | 12.4 | * | * | * | * | * | 2.6 | 7.8 | 18.7 | 46.9 | 111.1 | 298.8 |
| 1999..... | 11.8 | * | * | * | * | 0.7 | 2.6 | 7.1 | 18.9 | 43.0 | 112.6 | 279.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Cerebrovascular diseases (I60-I69) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 60.9 | 3.2 | 0.3 | 0.2 | 0.5 | 1.6 | 5.8 | 16.2 | 41.5 | 130.2 | 463.1 | 1,568.4 |
| 1999..... | 61.4 | 2.7 | 0.3 | 0.2 | 0.5 | 1.5 | 5.7 | 15.5 | 41.3 | 132.2 | 472.8 | 1,606.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 48.1 | 3.8 | * | 0.2 | 0.5 | 1.7 | 5.8 | 17.8 | 48.1 | 146.8 | 482.5 | 1,408.3 |
| 1999..... | 48.4 | 3.3 | 0.3 | 0.2 | 0.5 | 1.6 | 5.9 | 17.1 | 47.6 | 149.0 | 494.5 | 1,455.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 73.2 | 2.7 | 0.4 | 0.2 | 0.5 | 1.6 | 5.8 | 14.6 | 35.6 | 116.5 | 450.0 | 1,637.4 |
| 1999..... | 73.8 | 2.0 | 0.3 | 0.2 | 0.5 | 1.5 | 5.6 | 14.0 | 35.5 | 118.5 | 458.2 | 1,670.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 63.9 | 2.5 | 0.2 | 0.2 | 0.5 | 1.4 | 4.5 | 12.5 | 35.1 | 121.1 | 453.5 | 1,583.8 |
| 1999..... | 64.5 | 2.3 | 0.3 | 0.2 | 0.4 | 1.3 | 4.5 | 11.9 | 34.5 | 122.7 | 464.3 | 1,628.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 49.4 | 3.0 | * | 0.2 | 0.5 | 1.5 | 4.5 | 13.8 | 40.2 | 135.9 | 472.0 | 1,421.8 |
| 1999..... | 49.7 | 2.8 | * | 0.2 | 0.5 | 1.4 | 4.6 | 13.0 | 39.6 | 138.0 | 485.1 | 1,475.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 77.9 | 2.0 | * | 0.2 | 0.4 | 1.3 | 4.4 | 11.2 | 30.4 | 108.8 | 441.0 | 1,653.1 |
| 1999..... | 78.7 | 1.8 | * | 0.1 | 0.4 | 1.2 | 4.3 | 10.9 | 29.7 | 109.9 | 450.3 | 1,692.6 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 54.4 | 7.9 | * | * | 0.7 | 3.1 | 14.7 | 44.5 | 95.0 | 217.3 | 599.8 | 1,533.9 |
| 1999..... | 54.2 | 4.9 | * | * | 0.7 | 2.8 | 14.4 | 42.8 | 97.9 | 227.2 | 601.9 | 1,495.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Cerebrovascular diseases (I60-I69) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 47.8 | 9.1 | * | * | * | 2.9 | 14.8 | 51.6 | 120.5 | 256.6 | 640.1 | 1,346.2 |
| 1999..... | 47.7 | * | * | * | * | 2.9 | 14.8 | 51.0 | 123.3 | 264.0 | 634.9 | 1,343.1 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 60.4 | * | * | * | 0.8 | 3.2 | 14.5 | 38.5 | 75.5 | 188.7 | 575.4 | 1,612.4 |
| 1999..... | 60.0 | * | * | * | 0.8 | 2.8 | 14.1 | 36.0 | 78.5 | 200.3 | 582.1 | 1,559.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Atherosclerosis (I70) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.2 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.8 | 34.7 | 186.3 |
| 1999..... | 5.5 | * | * | * | * | * | 0.1 | 0.5 | 2.1 | 8.1 | 36.1 | 201.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 4.0 | * | * | * | * | * | 0.2 | 0.7 | 2.6 | 9.8 | 39.0 | 167.5 |
| 1999..... | 4.1 | * | * | * | * | * | 0.1 | 0.7 | 2.8 | 9.7 | 39.6 | 181.8 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 6.4 | * | * | * | * | * | * | 0.3 | 1.3 | 6.3 | 31.9 | 194.5 |
| 1999..... | 6.8 | * | * | * | * | * | * | 0.3 | 1.5 | 6.8 | 33.8 | 209.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.8 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 7.9 | 35.2 | 193.0 |
| 1999..... | 6.1 | * | * | * | * | * | 0.1 | 0.5 | 2.0 | 8.1 | 36.7 | 207.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 9.9 | 39.7 | 173.0 |
| 1999..... | 4.5 | * | * | * | * | * | 0.1 | 0.7 | 2.6 | 9.7 | 39.8 | 187.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 7.2 | * | * | * | * | * | * | 0.2 | 1.1 | 6.2 | 32.2 | 201.6 |
| 1999..... | 7.7 | * | * | * | * | * | * | 0.3 | 1.4 | 6.9 | 34.6 | 215.5 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.0 | * | * | * | * | * | * | 1.1 | 3.3 | 8.7 | 35.9 | 138.6 |
| 1999..... | 3.1 | * | * | * | * | * | * | 0.7 | 3.3 | 9.0 | 35.5 | 154.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Atherosclerosis (I70) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 2.5 | * | * | * | * | * | * | 1.5 | 4.1 | 10.5 | 39.1 | 136.2 |
| 1999..... | 2.6 | * | * | * | * | * | * | * | 4.6 | 11.3 | 42.9 | 152.1 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.4 | * | * | * | * | * | * | * | 2.6 | 7.4 | 33.9 | 139.6 |
| 1999..... | 3.5 | * | * | * | * | * | * | * | 2.3 | 7.3 | 31.1 | 155.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other diseases of circulatory system (I71-I78) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 9.5 | * | * | * | 0.2 | 0.4 | 1.0 | 2.8 | 9.3 | 32.6 | 74.9 | 161.1 |
| 1999..... | 9.5 | * | * | * | 0.1 | 0.4 | 1.1 | 2.7 | 9.4 | 31.9 | 76.3 | 164.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 10.5 | * | * | * | 0.2 | 0.6 | 1.2 | 3.8 | 13.4 | 46.0 | 101.1 | 203.9 |
| 1999..... | 10.3 | * | * | * | 0.2 | 0.5 | 1.3 | 3.8 | 13.6 | 44.9 | 103.0 | 199.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 8.5 | * | * | * | 0.1 | 0.3 | 0.8 | 1.8 | 5.6 | 21.6 | 57.2 | 142.6 |
| 1999..... | 8.6 | * | * | * | * | 0.2 | 0.8 | 1.7 | 5.6 | 21.2 | 58.4 | 149.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 10.1 | * | * | * | 0.1 | 0.4 | 0.8 | 2.4 | 8.9 | 33.2 | 75.3 | 161.2 |
| 1999..... | 10.2 | * | * | * | 0.1 | 0.3 | 1.0 | 2.4 | 9.3 | 32.5 | 76.9 | 164.8 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 11.3 | * | * | * | 0.2 | 0.5 | 1.0 | 3.4 | 13.0 | 47.4 | 102.9 | 209.5 |
| 1999..... | 11.3 | * | * | * | 0.2 | 0.5 | 1.2 | 3.4 | 13.6 | 46.4 | 105.0 | 204.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 9.0 | * | * | * | * | 0.2 | 0.6 | 1.4 | 5.1 | 21.3 | 56.5 | 140.6 |
| 1999..... | 9.2 | * | * | * | * | 0.2 | 0.7 | 1.5 | 5.2 | 20.9 | 57.9 | 148.2 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 7.4 | * | * | * | * | 0.7 | 2.2 | 5.7 | 14.6 | 32.2 | 79.4 | 178.1 |
| 1999..... | 7.0 | * | * | * | * | 0.6 | 2.0 | 5.4 | 12.5 | 31.0 | 78.1 | 179.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other diseases of circulatory system (I71-I78)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|------|------|------|-------|
| 2000..... | 7.2 | * | * | * | * | 0.9 | 2.9 | 7.8 | 19.3 | 39.3 | 92.9 | 163.9 |
| 1999..... | 6.7 | * | * | * | * | 0.9 | 2.3 | 7.6 | 16.5 | 37.9 | 87.6 | 170.6 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|-------|
| 2000..... | 7.5 | * | * | * | * | * | 1.6 | 3.9 | 11.0 | 27.0 | 71.3 | 184.1 |
| 1999..... | 7.2 | * | * | * | * | * | 1.7 | 3.6 | 9.5 | 26.1 | 72.3 | 182.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Aortic aneurysm and dissection (I71) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.7 | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 6.2 | 22.1 | 47.4 | 78.1 |
| 1999..... | 5.8 | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 | 6.1 | 22.0 | 48.9 | 79.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 7.2 | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 9.7 | 33.3 | 69.4 | 120.8 |
| 1999..... | 7.2 | * | * | * | 0.1 | 0.4 | 1.0 | 2.8 | 9.8 | 33.1 | 71.9 | 120.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | * | 0.2 | 0.4 | 0.9 | 3.0 | 12.9 | 32.5 | 59.8 |
| 1999..... | 4.4 | * | * | * | * | * | 0.4 | 0.8 | 2.9 | 12.9 | 33.5 | 62.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.3 | * | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 6.1 | 23.0 | 48.6 | 79.9 |
| 1999..... | 6.4 | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 6.2 | 22.9 | 50.4 | 82.1 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 7.9 | * | * | * | * | 0.4 | 0.8 | 2.5 | 9.6 | 35.0 | 71.8 | 126.6 |
| 1999..... | 8.0 | * | * | * | * | 0.4 | 0.9 | 2.4 | 10.0 | 34.9 | 75.0 | 125.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.7 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.8 | 12.9 | 32.8 | 59.9 |
| 1999..... | 4.8 | * | * | * | * | * | 0.4 | 0.7 | 2.7 | 12.9 | 33.9 | 63.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.6 | * | * | * | * | 0.5 | 1.6 | 3.3 | 8.3 | 16.6 | 37.0 | 62.4 |
| 1999..... | 3.4 | * | * | * | * | 0.4 | 1.4 | 3.9 | 6.6 | 16.3 | 34.4 | 60.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Aortic aneurysm and dissection (I71) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 4.0 | * | * | * | * | * | 2.4 | 4.8 | 12.0 | 21.1 | 45.6 | 63.9 |
| 1999..... | 3.7 | * | * | * | * | * | 2.0 | 6.1 | 9.6 | 21.1 | 37.5 | 70.6 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.3 | * | * | * | * | * | 0.9 | 2.0 | 5.5 | 13.2 | 31.8 | 61.8 |
| 1999..... | 3.1 | * | * | * | * | * | 0.9 | 2.0 | 4.4 | 12.9 | 32.6 | 56.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other diseases of arteries, arterioles and capillaries (I72-I78) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.7 | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 3.1 | 10.5 | 27.5 | 82.9 |
| 1999..... | 3.7 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.3 | 9.9 | 27.4 | 84.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 3.3 | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.7 | 12.7 | 31.8 | 83.1 |
| 1999..... | 3.1 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.9 | 11.8 | 31.1 | 79.1 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.1 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.6 | 8.8 | 24.6 | 82.9 |
| 1999..... | 4.2 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.8 | 8.4 | 24.9 | 86.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.8 | 10.2 | 26.7 | 81.4 |
| 1999..... | 3.8 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.1 | 9.6 | 26.5 | 82.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 3.4 | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.4 | 12.4 | 31.1 | 82.9 |
| 1999..... | 3.3 | * | * | * | * | * | 0.3 | 0.9 | 3.6 | 11.6 | 30.0 | 78.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.3 | * | * | * | * | * | 0.3 | 0.7 | 2.3 | 8.3 | 23.7 | 80.7 |
| 1999..... | 4.4 | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 8.0 | 24.0 | 84.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.7 | * | * | * | * | * | 0.6 | 2.3 | 6.3 | 15.6 | 42.4 | 115.7 |
| 1999..... | 3.6 | * | * | * | * | * | 0.6 | 1.5 | 5.9 | 14.7 | 43.6 | 118.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other diseases of arteries, arterioles and capillaries (I72-I78) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 3.2 | * | * | * | * | * | * | 2.9 | 7.3 | 18.2 | 47.3 | 100.0 |
| 1999..... | 3.0 | * | * | * | * | * | * | 1.5 | 6.9 | 16.8 | 50.1 | 100.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.2 | * | * | * | * | * | 0.7 | 1.9 | 5.5 | 13.7 | 39.5 | 122.3 |
| 1999..... | 4.2 | * | * | * | * | * | 0.8 | 1.6 | 5.2 | 13.2 | 39.8 | 126.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other disorders of circulatory system (I80-I99) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.7 | 0.7 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.2 | 4.4 | 9.4 | 25.2 |
| 1999..... | 1.5 | 0.6 | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 1.9 | 3.8 | 9.2 | 23.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | 0.3 | 0.9 | 1.6 | 2.3 | 4.8 | 9.5 | 21.1 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.0 | 4.3 | 9.5 | 20.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | 0.1 | 0.4 | 0.6 | 1.2 | 2.0 | 4.1 | 9.4 | 27.0 |
| 1999..... | 1.7 | * | * | * | * | 0.3 | 0.5 | 1.0 | 1.8 | 3.4 | 9.0 | 24.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.7 | 0.7 | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 1.9 | 4.0 | 9.2 | 25.2 |
| 1999..... | 1.5 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 3.6 | 8.9 | 23.1 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | 0.2 | 0.8 | 1.5 | 2.0 | 4.5 | 9.3 | 21.0 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.6 | 0.9 | 2.0 | 4.0 | 9.1 | 20.0 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | * | * | 0.3 | 0.5 | 1.0 | 1.8 | 3.6 | 9.0 | 27.0 |
| 1999..... | 1.7 | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.7 | 3.2 | 8.8 | 24.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | * | * | 0.9 | 1.5 | 2.9 | 4.5 | 8.7 | 14.7 | 30.1 |
| 1999..... | 1.8 | * | * | * | * | 0.5 | 1.4 | 2.4 | 2.9 | 6.5 | 14.1 | 28.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other disorders of circulatory system (I80-I99)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|-----|------|------|
| 2000..... | 2.0 | * | * | * | * | 0.9 | 1.8 | 3.5 | 5.4 | 8.7 | 12.9 | 27.7 |
| 1999..... | 1.7 | * | * | * | * | * | 1.8 | 3.0 | 2.8 | 7.9 | 16.5 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|-----|------|------|
| 2000..... | 2.3 | * | * | * | * | 1.0 | 1.3 | 2.3 | 3.9 | 8.7 | 15.7 | 31.1 |
| 1999..... | 1.8 | * | * | * | * | * | 1.1 | 2.0 | 3.1 | 5.5 | 12.6 | 31.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Influenza and pneumonia (J10-J18) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 23.7 | 7.5 | 0.7 | 0.2 | 0.5 | 1.0 | 2.4 | 4.8 | 12.0 | 39.6 | 161.0 | 734.4 |
| 1999..... | 23.4 | 8.4 | 0.9 | 0.2 | 0.5 | 0.9 | 2.4 | 4.7 | 11.2 | 37.7 | 158.0 | 748.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 21.3 | 8.3 | 0.7 | 0.2 | 0.5 | 1.1 | 2.8 | 5.9 | 14.7 | 48.9 | 196.5 | 846.3 |
| 1999..... | 20.8 | 8.8 | 0.9 | 0.3 | 0.5 | 1.0 | 2.7 | 6.1 | 13.1 | 47.5 | 194.2 | 860.8 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 26.1 | 6.7 | 0.7 | 0.2 | 0.5 | 0.8 | 1.9 | 3.7 | 9.6 | 31.9 | 137.0 | 686.2 |
| 1999..... | 25.8 | 8.0 | 0.9 | 0.2 | 0.5 | 0.7 | 2.1 | 3.5 | 9.5 | 29.6 | 133.8 | 700.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 25.6 | 5.7 | 0.6 | 0.2 | 0.5 | 0.9 | 2.1 | 4.1 | 11.0 | 37.6 | 160.4 | 749.9 |
| 1999..... | 25.2 | 6.1 | 0.6 | 0.2 | 0.4 | 0.7 | 1.9 | 4.0 | 10.2 | 36.5 | 158.7 | 762.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 22.5 | 6.3 | 0.5 | 0.2 | 0.5 | 0.9 | 2.5 | 4.9 | 13.1 | 45.8 | 194.5 | 864.2 |
| 1999..... | 22.0 | 6.6 | 0.5 | 0.3 | 0.4 | 0.9 | 2.1 | 5.0 | 11.7 | 45.4 | 194.1 | 877.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 28.6 | 5.0 | 0.7 | 0.2 | 0.4 | 0.8 | 1.7 | 3.4 | 9.0 | 30.7 | 137.2 | 701.0 |
| 1999..... | 28.4 | 5.5 | 0.7 | 0.2 | 0.4 | 0.6 | 1.8 | 3.1 | 8.8 | 29.2 | 134.9 | 714.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 17.0 | 17.7 | 1.3 | * | 0.8 | 1.8 | 4.5 | 10.3 | 22.3 | 58.7 | 174.3 | 602.5 |
| 1999..... | 16.9 | 20.2 | 1.9 | * | 0.9 | 2.0 | 5.5 | 11.0 | 21.2 | 53.0 | 160.4 | 636.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Influenza and pneumonia (J10-J18) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 17.4 | 18.9 | 1.9 | * | 0.7 | 2.4 | 5.7 | 14.5 | 31.3 | 79.9 | 227.3 | 697.0 |
| 1999..... | 17.1 | 20.1 | 2.2 | * | 0.8 | 2.3 | 6.7 | 15.8 | 28.5 | 73.9 | 208.5 | 745.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 16.6 | 16.4 | * | * | 0.8 | 1.2 | 3.5 | 6.8 | 15.4 | 43.4 | 142.1 | 562.9 |
| 1999..... | 16.7 | 20.4 | * | * | 0.9 | 1.8 | 4.5 | 6.9 | 15.7 | 37.7 | 131.5 | 590.6 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Pneumonia (J12-J18) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 23.1 | 7.3 | 0.6 | 0.2 | 0.5 | 0.9 | 2.3 | 4.6 | 11.6 | 38.5 | 157.0 | 714.5 |
| 1999..... | 22.8 | 8.0 | 0.8 | 0.2 | 0.4 | 0.9 | 2.3 | 4.7 | 10.9 | 36.8 | 154.3 | 726.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 20.8 | 8.0 | 0.6 | 0.2 | 0.5 | 1.1 | 2.8 | 5.7 | 14.3 | 47.6 | 192.5 | 826.7 |
| 1999..... | 20.3 | 8.5 | 0.7 | 0.2 | 0.5 | 1.0 | 2.6 | 6.0 | 12.8 | 46.6 | 190.0 | 841.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 25.3 | 6.5 | 0.6 | 0.2 | 0.4 | 0.8 | 1.9 | 3.6 | 9.2 | 30.9 | 133.0 | 666.1 |
| 1999..... | 25.1 | 7.6 | 0.8 | 0.2 | 0.4 | 0.7 | 2.1 | 3.4 | 9.2 | 28.7 | 130.4 | 678.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 24.9 | 5.4 | 0.5 | 0.2 | 0.4 | 0.8 | 2.0 | 4.0 | 10.5 | 36.5 | 156.2 | 728.8 |
| 1999..... | 24.5 | 5.8 | 0.6 | 0.2 | 0.4 | 0.7 | 1.9 | 4.0 | 9.9 | 35.6 | 154.8 | 740.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 21.9 | 6.0 | 0.4 | 0.2 | 0.5 | 0.9 | 2.4 | 4.8 | 12.7 | 44.4 | 190.4 | 843.6 |
| 1999..... | 21.5 | 6.6 | 0.4 | 0.2 | 0.4 | 0.8 | 2.1 | 4.9 | 11.3 | 44.5 | 189.7 | 856.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 27.7 | 4.9 | 0.6 | 0.2 | 0.4 | 0.8 | 1.7 | 3.3 | 8.5 | 29.8 | 132.9 | 679.7 |
| 1999..... | 27.5 | 5.1 | 0.7 | 0.2 | 0.3 | 0.6 | 1.7 | 3.0 | 8.6 | 28.3 | 131.3 | 691.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.7 | 17.3 | 1.3 | * | 0.7 | 1.8 | 4.5 | 10.1 | 22.1 | 57.8 | 172.9 | 593.1 |
| 1999..... | 16.7 | 19.5 | 1.9 | * | 0.8 | 2.0 | 5.4 | 10.9 | 21.0 | 52.2 | 158.7 | 630.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Pneumonia (J12-J18) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 17.2 | 18.9 | 1.9 | * | 0.7 | 2.3 | 5.6 | 14.2 | 31.0 | 79.2 | 225.8 | 685.3 |
| 1999..... | 16.9 | 19.0 | 2.1 | * | 0.8 | 2.3 | 6.6 | 15.7 | 28.4 | 73.1 | 206.7 | 742.2 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 16.3 | 15.7 | * | * | 0.8 | 1.2 | 3.5 | 6.7 | 15.2 | 42.2 | 140.7 | 554.5 |
| 1999..... | 16.4 | 20.0 | * | * | 0.9 | 1.7 | 4.4 | 6.9 | 15.4 | 37.0 | 129.9 | 583.8 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

BLACK, FEMALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Chronic lower respiratory diseases (J40-J47) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 44.3 | 0.9 | 0.3 | 0.4 | 0.5 | 0.8 | 2.1 | 8.8 | 44.8 | 171.5 | 387.6 | 640.1 |
| 1999..... | 45.5 | 0.9 | 0.4 | 0.4 | 0.6 | 0.9 | 2.0 | 8.7 | 48.3 | 179.2 | 400.4 | 642.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 44.6 | 1.2 | 0.4 | 0.4 | 0.6 | 0.9 | 2.0 | 9.1 | 48.7 | 197.6 | 480.3 | 918.3 |
| 1999..... | 46.8 | * | 0.4 | 0.5 | 0.6 | 0.8 | 1.8 | 8.9 | 53.4 | 213.2 | 507.2 | 959.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 44.1 | * | 0.3 | 0.3 | 0.4 | 0.7 | 2.2 | 8.4 | 41.3 | 150.0 | 324.9 | 520.2 |
| 1999..... | 44.3 | * | 0.3 | 0.2 | 0.5 | 0.9 | 2.1 | 8.5 | 43.6 | 151.4 | 328.9 | 509.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 49.9 | 0.8 | 0.2 | 0.2 | 0.3 | 0.6 | 1.7 | 8.3 | 46.3 | 181.2 | 405.4 | 668.1 |
| 1999..... | 51.1 | * | 0.3 | 0.2 | 0.4 | 0.6 | 1.6 | 8.3 | 49.9 | 188.7 | 416.7 | 671.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 49.3 | * | * | 0.2 | 0.4 | 0.6 | 1.6 | 8.5 | 49.2 | 204.4 | 495.2 | 951.4 |
| 1999..... | 51.7 | * | * | 0.3 | 0.4 | 0.5 | 1.5 | 8.3 | 54.0 | 220.5 | 519.8 | 997.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 50.4 | * | * | 0.2 | 0.3 | 0.6 | 1.8 | 8.0 | 43.5 | 161.8 | 344.4 | 547.0 |
| 1999..... | 50.5 | * | * | 0.2 | 0.4 | 0.6 | 1.8 | 8.2 | 46.1 | 162.2 | 347.3 | 535.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 21.5 | * | 0.9 | 1.1 | 1.4 | 2.2 | 4.8 | 14.5 | 44.5 | 118.6 | 239.1 | 374.8 |
| 1999..... | 22.7 | * | * | 1.1 | 1.4 | 2.7 | 4.7 | 14.3 | 46.5 | 125.8 | 267.7 | 377.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Chronic lower respiratory diseases (J40-J47)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|-----|-----|------|------|-------|-------|-------|
| 2000..... | 25.3 | * | * | 1.4 | 1.7 | 2.9 | 4.8 | 15.7 | 57.0 | 169.1 | 359.6 | 639.6 |
| 1999..... | 27.2 | * | * | 1.5 | 1.8 | 2.8 | 4.7 | 16.0 | 62.0 | 177.4 | 423.9 | 650.9 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|-----|-----|-----|------|------|------|-------|-------|
| 2000..... | 18.2 | * | * | 0.9 | 1.0 | 1.6 | 4.8 | 13.5 | 34.9 | 81.9 | 165.9 | 264.1 |
| 1999..... | 18.7 | * | * | 0.7 | 0.9 | 2.7 | 4.6 | 12.8 | 34.7 | 88.3 | 174.0 | 262.9 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

| ALL RACES 3/, MALE | | | | | | | | | | | | |
|--------------------|-----|---|---|---|---|---|---|---|-----|-----|-----|------|
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.5 | 12.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.5 | 1.1 | 3.2 | 11.4 |

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|-----|------|
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 11.4 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.5 | 10.4 |

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|-----|------|
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 12.5 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.8 | 11.2 |

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|-----|-----|------|
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.7 | 13.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.5 | 1.2 | 3.1 | 12.0 |

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|-----|-----|------|
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.2 | 12.3 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.6 | 10.9 |

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Emphysema (J43) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.1 | * | * | * | * | * | 0.2 | 1.3 | 7.5 | 28.2 | 52.9 | 62.5 |
| 1999..... | 6.5 | * | * | * | * | * | 0.2 | 1.3 | 8.6 | 31.0 | 57.3 | 61.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.4 | * | * | * | * | * | 0.2 | 1.5 | 8.2 | 33.4 | 68.0 | 98.8 |
| 1999..... | 7.1 | * | * | * | * | * | 0.2 | 1.5 | 9.7 | 38.2 | 75.5 | 101.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 5.7 | * | * | * | * | * | 0.2 | 1.0 | 6.9 | 23.9 | 42.7 | 46.9 |
| 1999..... | 6.0 | * | * | * | * | * | 0.2 | 1.1 | 7.6 | 25.2 | 45.1 | 45.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.9 | * | * | * | * | * | 0.2 | 1.2 | 8.0 | 30.1 | 55.8 | 65.1 |
| 1999..... | 7.4 | * | * | * | * | * | 0.2 | 1.3 | 9.2 | 33.3 | 60.2 | 64.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 7.2 | * | * | * | * | * | 0.2 | 1.4 | 8.5 | 34.8 | 70.9 | 100.7 |
| 1999..... | 7.9 | * | * | * | * | * | 0.2 | 1.4 | 10.0 | 40.0 | 78.1 | 105.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 6.6 | * | * | * | * | * | 0.2 | 1.0 | 7.5 | 26.2 | 45.4 | 49.8 |
| 1999..... | 6.9 | * | * | * | * | * | 0.2 | 1.2 | 8.4 | 27.8 | 48.1 | 47.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.6 | * | * | * | * | * | * | 2.1 | 6.1 | 16.6 | 27.4 | 37.3 |
| 1999..... | 2.7 | * | * | * | * | * | 0.4 | 1.9 | 6.2 | 16.7 | 31.9 | 37.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Emphysema (J43) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 3.3 | * | * | * | * | * | * | 3.0 | 8.4 | 26.2 | 40.3 | 80.9 |
| 1999..... | 3.6 | * | * | * | * | * | * | 3.0 | 9.3 | 27.8 | 53.4 | 67.4 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.8 | * | * | * | * | * | * | 1.3 | 4.3 | 9.6 | 19.6 | 19.1 |
| 1999..... | 1.8 | * | * | * | * | * | * | 1.0 | 3.9 | 8.7 | 19.1 | 25.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Asthma (J45-J46) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.6 | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.0 | 1.7 | 2.3 | 4.0 | 6.9 | 16.5 |
| 1999..... | 1.7 | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.0 | 1.7 | 2.5 | 4.2 | 7.6 | 17.2 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.2 | * | 0.3 | 0.4 | 0.5 | 0.8 | 0.9 | 1.3 | 1.7 | 3.1 | 4.9 | 12.2 |
| 1999..... | 1.2 | * | * | 0.4 | 0.6 | 0.7 | 0.8 | 1.1 | 1.9 | 3.3 | 5.6 | 12.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.0 | * | * | 0.3 | 0.3 | 0.6 | 1.2 | 2.0 | 2.7 | 4.8 | 8.2 | 18.3 |
| 1999..... | 2.2 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.2 | 3.1 | 5.1 | 9.0 | 19.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.1 | 1.7 | 3.4 | 6.5 | 15.7 |
| 1999..... | 1.5 | * | * | 0.2 | 0.4 | 0.4 | 0.7 | 1.2 | 1.9 | 3.6 | 7.3 | 16.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 1.0 | 1.2 | 2.6 | 4.6 | 10.8 |
| 1999..... | 0.9 | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.8 | 1.4 | 2.5 | 5.2 | 10.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.8 | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.3 | 2.1 | 4.2 | 7.8 | 17.8 |
| 1999..... | 2.0 | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 1.7 | 2.4 | 4.4 | 8.7 | 19.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.3 | * | * | 1.1 | 1.3 | 2.0 | 3.1 | 5.5 | 7.0 | 9.0 | 10.6 | 22.3 |
| 1999..... | 3.3 | * | * | 1.1 | 1.2 | 2.5 | 3.0 | 5.4 | 6.9 | 9.4 | 10.6 | 20.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Asthma (J45-J46) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 2.9 | * | * | 1.3 | 1.5 | 2.6 | 2.9 | 3.9 | 6.4 | 7.6 | 6.2 | 23.4 |
| 1999..... | 2.9 | * | * | 1.5 | 1.6 | 2.7 | 2.8 | 4.0 | 5.6 | 8.8 | 8.1 | * |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.7 | * | * | 0.9 | 1.0 | 1.5 | 3.3 | 6.8 | 7.4 | 10.1 | 13.2 | 21.8 |
| 1999..... | 3.6 | * | * | 0.7 | 0.9 | 2.2 | 3.1 | 6.6 | 7.9 | 9.9 | 12.1 | 21.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other chronic lower respiratory diseases (J44,J47) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 36.2 | * | * | * | * | 0.1 | 0.8 | 5.7 | 34.7 | 138.4 | 325.1 | 549.5 |
| 1999..... | 36.9 | * | * | * | * | 0.1 | 0.7 | 5.6 | 36.8 | 142.9 | 332.7 | 553.0 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 36.5 | * | * | * | * | * | 0.8 | 6.2 | 38.4 | 160.2 | 403.9 | 795.1 |
| 1999..... | 38.1 | * | * | * | * | * | 0.8 | 6.1 | 41.4 | 170.5 | 422.9 | 833.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 35.9 | * | * | * | * | * | 0.8 | 5.3 | 31.4 | 120.5 | 271.8 | 443.7 |
| 1999..... | 35.7 | * | * | * | * | 0.2 | 0.7 | 5.1 | 32.5 | 120.4 | 272.4 | 434.3 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 41.1 | * | * | * | * | * | 0.8 | 5.8 | 36.3 | 146.8 | 340.3 | 574.8 |
| 1999..... | 41.7 | * | * | * | * | 0.1 | 0.7 | 5.7 | 38.4 | 150.8 | 346.5 | 579.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 40.8 | * | * | * | * | * | 0.8 | 6.1 | 39.2 | 166.0 | 416.0 | 826.7 |
| 1999..... | 42.4 | * | * | * | * | * | 0.7 | 6.0 | 42.1 | 176.8 | 433.3 | 869.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 41.5 | * | * | * | * | * | 0.8 | 5.5 | 33.6 | 130.6 | 288.9 | 467.1 |
| 1999..... | 41.1 | * | * | * | * | 0.1 | 0.7 | 5.3 | 34.9 | 129.2 | 288.0 | 457.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 15.5 | * | * | * | * | * | 1.3 | 6.8 | 31.3 | 92.1 | 198.9 | 311.4 |
| 1999..... | 16.5 | * | * | * | * | * | 1.2 | 6.6 | 33.0 | 98.8 | 222.3 | 314.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other chronic lower respiratory diseases (J44,J47)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|---|-----|-----|------|-------|-------|-------|
| 2000..... | 18.9 | * | * | * | * | * | 1.3 | 8.7 | 41.9 | 134.5 | 311.1 | 532.1 |
| 1999..... | 20.4 | * | * | * | * | * | 1.5 | 8.6 | 46.6 | 139.8 | 357.6 | 562.9 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|---|-----|-----|------|------|-------|-------|
| 2000..... | 12.4 | * | * | * | * | * | 1.3 | 5.2 | 23.2 | 61.3 | 130.7 | 219.2 |
| 1999..... | 13.0 | * | * | * | * | * | 1.0 | 5.0 | 22.7 | 68.9 | 141.1 | 210.9 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Pneumonitis due to solids and liquids (J69) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.0 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 10.5 | 44.7 | 185.2 |
| 1999..... | 5.6 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.5 | 9.6 | 41.4 | 174.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.2 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.1 | 14.6 | 61.2 | 261.9 |
| 1999..... | 5.8 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 3.3 | 13.2 | 59.0 | 248.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 5.9 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.9 | 7.1 | 33.6 | 152.1 |
| 1999..... | 5.4 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.7 | 29.6 | 143.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.6 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.4 | 10.0 | 44.6 | 188.9 |
| 1999..... | 6.0 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.3 | 9.2 | 41.0 | 177.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 6.7 | * | * | * | * | 0.1 | 0.5 | 1.1 | 2.9 | 13.6 | 61.1 | 270.9 |
| 1999..... | 6.2 | * | * | * | * | 0.2 | 0.4 | 0.9 | 3.0 | 12.4 | 58.0 | 255.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 6.4 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.9 | 33.3 | 153.8 |
| 1999..... | 5.9 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.7 | 6.5 | 29.5 | 145.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.3 | * | * | * | * | * | 0.7 | 1.9 | 4.2 | 17.1 | 52.4 | 160.3 |
| 1999..... | 4.3 | * | * | * | * | * | 0.7 | 1.5 | 4.5 | 15.8 | 53.2 | 167.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Pneumonitis due to solids and liquids (J69)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|------|------|-------|
| 2000..... | 4.6 | * | * | * | * | * | 0.8 | 2.8 | 5.3 | 26.6 | 72.3 | 193.7 |
| 1999..... | 4.7 | * | * | * | * | * | 0.9 | 1.9 | 6.7 | 24.0 | 78.9 | 211.9 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|------|------|-------|
| 2000..... | 4.1 | * | * | * | * | * | * | 1.2 | 3.4 | 10.2 | 40.4 | 146.3 |
| 1999..... | 4.0 | * | * | * | * | * | * | 1.2 | 2.8 | 9.9 | 37.8 | 148.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 9.3 | 7.1 | 0.6 | 0.2 | 0.3 | 0.7 | 1.4 | 3.6 | 10.1 | 30.0 | 72.3 | 139.9 |
| 1999..... | 9.2 | 6.6 | 0.6 | 0.2 | 0.3 | 0.7 | 1.3 | 3.4 | 9.7 | 29.7 | 71.2 | 146.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 9.2 | 7.8 | 0.6 | 0.2 | 0.4 | 0.7 | 1.5 | 3.9 | 11.6 | 36.1 | 89.0 | 170.5 |
| 1999..... | 9.1 | 7.4 | 0.7 | 0.2 | 0.4 | 0.7 | 1.4 | 3.6 | 10.9 | 36.7 | 88.8 | 175.4 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 9.3 | 6.3 | 0.6 | 0.2 | 0.3 | 0.7 | 1.2 | 3.3 | 8.7 | 25.0 | 61.0 | 126.7 |
| 1999..... | 9.3 | 5.7 | 0.5 | 0.2 | 0.2 | 0.7 | 1.1 | 3.2 | 8.6 | 24.0 | 59.3 | 134.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 10.0 | 5.5 | 0.5 | 0.2 | 0.3 | 0.6 | 1.2 | 3.2 | 9.8 | 30.5 | 73.9 | 144.3 |
| 1999..... | 9.9 | 4.9 | 0.5 | 0.2 | 0.3 | 0.6 | 1.1 | 3.0 | 9.4 | 30.0 | 72.7 | 150.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 9.9 | 6.4 | 0.5 | 0.2 | 0.3 | 0.6 | 1.3 | 3.5 | 11.2 | 36.6 | 91.0 | 175.9 |
| 1999..... | 9.8 | 5.5 | 0.6 | 0.2 | 0.4 | 0.6 | 1.2 | 3.3 | 10.4 | 37.4 | 91.0 | 179.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.2 | 4.5 | 0.6 | 0.2 | 0.3 | 0.6 | 1.1 | 3.0 | 8.4 | 25.3 | 62.3 | 130.8 |
| 1999..... | 10.0 | 4.2 | 0.5 | 0.2 | 0.2 | 0.5 | 0.9 | 2.8 | 8.3 | 23.9 | 60.3 | 138.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.7 | 17.7 | 0.9 | * | 0.7 | 1.1 | 2.7 | 7.0 | 13.6 | 29.7 | 59.1 | 98.5 |
| 1999..... | 6.8 | 16.4 | 1.2 | * | 0.4 | 1.4 | 2.9 | 6.6 | 13.6 | 30.2 | 59.0 | 111.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|------|---|---|-----|-----|-----|-----|------|------|------|-------|
| 2000..... | 6.8 | 17.2 | * | * | 0.8 | 1.2 | 3.2 | 8.1 | 15.9 | 34.4 | 70.9 | 122.4 |
| 1999..... | 6.6 | 18.7 | * | * | * | 1.1 | 2.9 | 7.5 | 16.3 | 33.8 | 70.8 | 134.7 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|---|---|---|-----|-----|-----|------|------|------|-------|
| 2000..... | 6.6 | 18.2 | * | * | * | 1.1 | 2.2 | 6.0 | 11.8 | 26.3 | 52.0 | 88.5 |
| 1999..... | 6.9 | 13.9 | * | * | * | 1.7 | 2.9 | 5.9 | 11.6 | 27.6 | 52.0 | 102.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Peptic ulcer (K25-K28) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.7 | * | * | * | * | * | 0.3 | 0.8 | 1.8 | 4.1 | 11.5 | 35.1 |
| 1999..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 4.6 | 12.3 | 34.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.6 | * | * | * | * | * | 0.4 | 1.2 | 2.5 | 5.4 | 13.4 | 38.6 |
| 1999..... | 1.6 | * | * | * | * | * | 0.3 | 1.0 | 2.4 | 6.1 | 14.6 | 36.1 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.7 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.1 | 10.2 | 33.6 |
| 1999..... | 1.7 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.3 | 10.8 | 33.8 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.8 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.1 | 11.6 | 35.9 |
| 1999..... | 1.8 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.5 | 12.4 | 35.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.7 | * | * | * | * | * | 0.3 | 1.1 | 2.4 | 5.3 | 13.3 | 38.8 |
| 1999..... | 1.7 | * | * | * | * | * | 0.2 | 0.8 | 2.2 | 6.0 | 14.7 | 37.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.8 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.1 | 10.4 | 34.6 |
| 1999..... | 1.9 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.3 | 10.8 | 34.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | * | * | * | 0.5 | 1.7 | 2.7 | 4.7 | 10.2 | 26.7 |
| 1999..... | 1.3 | * | * | * | * | * | 0.8 | 1.5 | 2.8 | 5.5 | 11.1 | 25.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Peptic ulcer (K25-K28)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|------|------|
| 2000..... | 1.4 | * | * | * | * | * | 0.8 | 2.4 | 3.8 | 6.7 | 13.5 | 28.7 |
| 1999..... | 1.5 | * | * | * | * | * | 1.2 | 2.4 | 4.6 | 7.8 | 12.0 | 26.1 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|------|------|
| 2000..... | 1.1 | * | * | * | * | * | * | 1.1 | 1.8 | 3.3 | 8.2 | 25.8 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 3.9 | 10.6 | 25.3 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEALL RACES 3/, FEMALEWHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Hernia (K40-K46) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.6 | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.8 | 12.9 |
| 1999..... | 0.5 | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.7 | 12.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.4 | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.9 | 11.9 |
| 1999..... | 0.4 | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.9 | 12.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.7 | 13.4 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.6 | 12.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.6 | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.8 | 13.2 |
| 1999..... | 0.6 | 1.1 | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.1 | 3.7 | 13.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.5 | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.0 | 12.1 |
| 1999..... | 0.5 | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 4.0 | 12.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.6 | 13.6 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 0.4 | 1.1 | 3.5 | 13.2 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 1.2 | 4.2 | 12.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 1.4 | 4.3 | 9.9 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Chronic liver disease and cirrhosis (K70,K73-K74) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 9.6 | * | * | * | 0.1 | 1.1 | 7.5 | 17.9 | 24.1 | 30.2 | 31.1 | 22.8 |
| 1999..... | 9.6 | * | * | * | 0.1 | 1.1 | 7.4 | 17.8 | 24.1 | 31.0 | 32.1 | 23.1 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 12.8 | * | * | * | * | 1.5 | 10.4 | 27.4 | 35.0 | 40.4 | 37.8 | 32.6 |
| 1999..... | 12.8 | * | * | * | * | 1.4 | 10.4 | 27.6 | 35.4 | 42.1 | 39.1 | 31.8 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 6.6 | * | * | * | * | 0.7 | 4.7 | 8.8 | 14.1 | 21.7 | 26.6 | 18.6 |
| 1999..... | 6.6 | * | * | * | * | 0.8 | 4.4 | 8.4 | 13.8 | 21.8 | 27.4 | 19.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 10.2 | * | * | * | 0.1 | 1.0 | 7.4 | 17.6 | 24.0 | 31.0 | 32.3 | 23.7 |
| 1999..... | 10.1 | * | * | * | * | 1.1 | 7.2 | 17.3 | 23.6 | 31.8 | 33.2 | 23.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 13.5 | * | * | * | * | 1.5 | 10.4 | 27.1 | 34.9 | 41.6 | 39.2 | 33.9 |
| 1999..... | 13.4 | * | * | * | * | 1.4 | 10.3 | 26.9 | 34.7 | 43.0 | 40.3 | 33.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 6.9 | * | * | * | * | 0.6 | 4.5 | 8.3 | 13.8 | 22.2 | 27.7 | 19.3 |
| 1999..... | 6.9 | * | * | * | * | 0.7 | 4.2 | 7.9 | 13.3 | 22.5 | 28.4 | 19.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 7.8 | * | * | * | * | 1.1 | 7.9 | 22.7 | 26.2 | 25.6 | 18.3 | 13.8 |
| 1999..... | 8.1 | * | * | * | * | 1.0 | 8.1 | 23.5 | 29.8 | 26.1 | 21.9 | 17.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Chronic liver disease and cirrhosis (K70,K73-K74)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|---|---|-----|------|------|------|------|------|---|
| 2000..... | 10.3 | * | * | * | * | 1.1 | 10.6 | 34.0 | 39.7 | 36.1 | 23.8 | * |
| 1999..... | 11.1 | * | * | * | * | 1.1 | 11.2 | 37.2 | 45.3 | 39.2 | 28.8 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|------|------|------|------|------|
| 2000..... | 5.4 | * | * | * | * | 1.1 | 5.5 | 13.2 | 16.0 | 17.9 | 15.0 | 12.0 |
| 1999..... | 5.4 | * | * | * | * | 1.0 | 5.4 | 12.1 | 18.0 | 16.6 | 17.8 | 16.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Alcoholic liver disease (K70) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | * | 0.7 | 5.0 | 10.5 | 12.4 | 10.4 | 6.1 | 2.2 |
| 1999..... | 4.4 | * | * | * | * | 0.7 | 4.9 | 10.6 | 12.2 | 10.6 | 6.5 | 2.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.7 | * | * | * | * | 1.1 | 6.9 | 16.3 | 19.4 | 17.5 | 11.3 | 5.0 |
| 1999..... | 6.7 | * | * | * | * | 0.9 | 6.8 | 16.7 | 19.7 | 18.3 | 11.0 | 5.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 3.1 | 4.8 | 6.0 | 4.6 | 2.6 | 1.0 |
| 1999..... | 2.2 | * | * | * | * | 0.5 | 3.0 | 4.8 | 5.4 | 4.4 | 3.4 | 1.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.5 | * | * | * | * | 0.7 | 4.9 | 10.4 | 12.3 | 10.6 | 6.1 | 2.3 |
| 1999..... | 4.5 | * | * | * | * | 0.7 | 4.8 | 10.4 | 11.9 | 10.6 | 6.5 | 2.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 7.0 | * | * | * | * | 1.1 | 6.8 | 16.4 | 19.4 | 17.8 | 11.5 | 5.2 |
| 1999..... | 6.9 | * | * | * | * | 0.9 | 6.7 | 16.5 | 19.3 | 18.2 | 11.0 | 5.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | * | * | 0.3 | 3.0 | 4.5 | 5.7 | 4.6 | 2.5 | 1.0 |
| 1999..... | 2.1 | * | * | * | * | 0.5 | 2.8 | 4.5 | 5.1 | 4.4 | 3.4 | 1.5 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.9 | * | * | * | * | 0.6 | 5.0 | 12.3 | 13.5 | 10.0 | 5.6 | * |
| 1999..... | 4.2 | * | * | * | * | 0.5 | 5.0 | 13.2 | 15.4 | 11.6 | 7.2 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Alcoholic liver disease (K70)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|------|------|------|---|
| 2000..... | 5.5 | * | * | * | * | * | 7.0 | 18.2 | 20.2 | 17.3 | 9.7 | * |
| 1999..... | 6.1 | * | * | * | * | * | 6.7 | 20.6 | 25.3 | 20.9 | 12.3 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|---|
| 2000..... | 2.5 | * | * | * | * | * | 3.3 | 7.4 | 8.3 | 4.7 | * | * |
| 1999..... | 2.4 | * | * | * | * | * | 3.5 | 7.0 | 7.8 | 4.7 | 4.1 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other chronic liver disease and cirrhosis (K73-K74) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.2 | * | * | * | 0.1 | 0.4 | 2.6 | 7.4 | 11.7 | 19.8 | 25.0 | 20.6 |
| 1999..... | 5.2 | * | * | * | * | 0.4 | 2.5 | 7.2 | 11.9 | 20.3 | 25.6 | 20.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.1 | * | * | * | * | 0.4 | 3.5 | 11.0 | 15.6 | 23.0 | 26.5 | 27.5 |
| 1999..... | 6.1 | * | * | * | * | 0.5 | 3.6 | 10.9 | 15.7 | 23.9 | 28.1 | 26.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | * | 0.3 | 1.6 | 4.0 | 8.1 | 17.1 | 24.0 | 17.6 |
| 1999..... | 4.4 | * | * | * | * | 0.3 | 1.4 | 3.6 | 8.4 | 17.5 | 24.0 | 17.9 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.6 | * | * | * | * | 0.3 | 2.5 | 7.2 | 11.7 | 20.4 | 26.2 | 21.4 |
| 1999..... | 5.6 | * | * | * | * | 0.4 | 2.5 | 6.9 | 11.7 | 21.1 | 26.7 | 21.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 6.5 | * | * | * | * | 0.4 | 3.5 | 10.8 | 15.5 | 23.7 | 27.7 | 28.7 |
| 1999..... | 6.5 | * | * | * | * | 0.5 | 3.5 | 10.4 | 15.4 | 24.8 | 29.3 | 27.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.8 | * | * | * | * | 0.3 | 1.5 | 3.8 | 8.2 | 17.6 | 25.2 | 18.2 |
| 1999..... | 4.7 | * | * | * | * | 0.3 | 1.4 | 3.4 | 8.2 | 18.1 | 25.0 | 18.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.8 | * | * | * | * | 0.4 | 2.9 | 10.4 | 12.8 | 15.6 | 12.8 | 12.2 |
| 1999..... | 3.9 | * | * | * | * | 0.5 | 3.1 | 10.3 | 14.4 | 14.5 | 14.7 | 12.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other chronic liver disease and cirrhosis (K73-K74)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|------|------|------|------|---|
| 2000..... | 4.8 | * | * | * | * | * | 3.6 | 15.9 | 19.5 | 18.7 | 14.1 | * |
| 1999..... | 5.0 | * | * | * | * | * | 4.4 | 16.5 | 20.0 | 18.2 | 16.5 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|------|------|------|------|
| 2000..... | 2.9 | * | * | * | * | * | 2.2 | 5.8 | 7.7 | 13.2 | 12.0 | 11.6 |
| 1999..... | 3.0 | * | * | * | * | * | 1.9 | 5.0 | 10.1 | 11.8 | 13.7 | 13.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Cholelithiasis and other disorders of gallbladder (K80-K82) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 7.5 | 24.5 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.6 | 7.8 | 23.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.9 | 9.3 | 26.2 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.1 | 9.2 | 26.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.1 | 6.3 | 23.8 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.2 | 6.9 | 22.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.3 | 7.4 | 25.3 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 7.9 | 24.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 9.4 | 27.5 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.0 | 9.5 | 27.2 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 6.1 | 24.4 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.3 | 0.6 | 2.1 | 6.8 | 23.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.7 | 1.2 | 3.8 | 8.4 | 17.9 |
| 1999..... | 0.7 | * | * | * | * | * | 0.4 | 0.5 | 1.5 | 3.5 | 7.3 | 15.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Cholelithiasis and other disorders of gallbladder (K80-K82)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|-----|-----|---|
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 3.5 | 7.9 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 3.5 | 6.9 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|-----|-----|------|
| 2000..... | 1.0 | * | * | * | * | * | * | * | * | 4.1 | 8.8 | 20.0 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 3.5 | 7.6 | 15.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 13.5 | 4.2 | * | 0.1 | 0.2 | 0.7 | 1.6 | 4.4 | 12.9 | 38.5 | 101.2 | 274.2 |
| 1999..... | 13.0 | 4.3 | * | 0.1 | 0.2 | 0.7 | 1.6 | 4.1 | 12.2 | 37.6 | 98.2 | 267.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 13.2 | 5.2 | * | * | 0.2 | 0.7 | 1.9 | 5.4 | 14.4 | 43.4 | 125.2 | 361.2 |
| 1999..... | 12.8 | 4.6 | * | * | 0.2 | 0.7 | 1.8 | 4.8 | 13.7 | 42.7 | 123.8 | 357.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 13.8 | 3.1 | * | * | 0.2 | 0.6 | 1.4 | 3.5 | 11.6 | 34.4 | 85.0 | 236.7 |
| 1999..... | 13.3 | 4.1 | * | * | 0.2 | 0.7 | 1.4 | 3.5 | 10.9 | 33.3 | 81.0 | 229.7 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 13.1 | 2.9 | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.0 | 9.7 | 32.4 | 94.3 | 265.7 |
| 1999..... | 12.5 | 3.4 | * | 0.1 | 0.1 | 0.5 | 1.0 | 2.7 | 9.3 | 31.9 | 90.6 | 255.2 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 12.9 | 3.7 | * | * | 0.2 | 0.5 | 1.3 | 3.6 | 11.1 | 38.3 | 119.1 | 358.8 |
| 1999..... | 12.4 | 4.0 | * | * | * | 0.5 | 1.2 | 3.2 | 10.7 | 37.4 | 116.0 | 349.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 13.2 | 2.0 | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 8.5 | 27.5 | 77.4 | 225.9 |
| 1999..... | 12.6 | 2.7 | * | * | 0.2 | 0.4 | 0.9 | 2.3 | 7.9 | 27.3 | 73.5 | 215.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 19.6 | 12.2 | * | * | 0.4 | 1.9 | 5.4 | 16.2 | 41.1 | 97.4 | 198.8 | 407.4 |
| 1999..... | 19.3 | 10.2 | * | * | 0.5 | 1.9 | 5.5 | 15.7 | 37.8 | 93.5 | 199.0 | 434.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|------|---|---|---|-----|-----|------|------|-------|-------|-------|
| 2000..... | 18.3 | 14.2 | * | * | * | 2.0 | 6.5 | 21.2 | 46.9 | 100.1 | 224.9 | 452.3 |
| 1999..... | 18.2 | 8.6 | * | * | * | 1.9 | 6.5 | 19.8 | 42.3 | 98.9 | 243.9 | 487.9 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|---|---|---|-----|-----|------|------|------|-------|-------|
| 2000..... | 20.7 | 10.1 | * | * | * | 1.8 | 4.3 | 12.0 | 36.7 | 95.4 | 182.9 | 388.6 |
| 1999..... | 20.2 | 11.8 | * | * | * | 2.0 | 4.6 | 12.3 | 34.4 | 89.5 | 172.0 | 411.8 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

ALL RACES 3/, BOTH SEXES

[illegible][illegible]

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|-----|-----|-----|
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 |

[illegible]

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|---|---|-----|-----|
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | * |

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07, N26)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|-----|
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.4 | 2.7 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.6 | 3.2 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|-----|
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.7 | 4.3 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 2.2 | 4.8 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|-----|-----|-----|
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.5 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|-----|-----|-----|-----|-----|-----|
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.4 | 2.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 2.9 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|-----|-----|-----|
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 | 4.5 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|---|-----|-----|-----|-----|
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 1.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.2 | 2.2 |

BLACK, BOTH SEXES

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Renal failure (N17-N19) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 13.2 | 4.0 | * | * | 0.2 | 0.6 | 1.6 | 4.3 | 12.6 | 37.7 | 99.3 | 270.1 |
| 1999..... | 12.7 | 4.3 | * | 0.1 | 0.2 | 0.6 | 1.5 | 4.0 | 11.9 | 36.8 | 95.9 | 263.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 12.9 | 5.1 | * | * | 0.2 | 0.6 | 1.8 | 5.2 | 14.1 | 42.6 | 122.9 | 355.0 |
| 1999..... | 12.5 | 4.5 | * | * | 0.1 | 0.6 | 1.7 | 4.7 | 13.4 | 42.0 | 120.9 | 350.8 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 13.5 | 2.8 | * | * | 0.2 | 0.6 | 1.3 | 3.4 | 11.3 | 33.7 | 83.3 | 233.5 |
| 1999..... | 13.0 | 4.1 | * | * | 0.2 | 0.6 | 1.3 | 3.3 | 10.6 | 32.5 | 79.2 | 226.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 12.8 | 2.6 | * | * | 0.2 | 0.4 | 1.1 | 2.9 | 9.5 | 31.7 | 92.4 | 262.0 |
| 1999..... | 12.2 | 3.3 | * | * | 0.1 | 0.4 | 1.0 | 2.6 | 9.0 | 31.2 | 88.5 | 251.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 12.7 | 3.5 | * | * | 0.2 | 0.4 | 1.2 | 3.4 | 10.8 | 37.4 | 116.9 | 353.3 |
| 1999..... | 12.1 | 3.8 | * | * | * | 0.4 | 1.1 | 3.0 | 10.5 | 36.7 | 113.2 | 343.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 13.0 | 1.7 | * | * | 0.2 | 0.4 | 0.9 | 2.3 | 8.2 | 26.8 | 75.8 | 223.0 |
| 1999..... | 12.3 | 2.7 | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 7.7 | 26.5 | 71.8 | 212.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 19.2 | 11.7 | * | * | 0.4 | 1.8 | 5.1 | 15.6 | 40.6 | 96.1 | 196.0 | 401.4 |
| 1999..... | 18.9 | 10.2 | * | * | 0.5 | 1.9 | 5.3 | 15.3 | 37.0 | 92.2 | 195.5 | 426.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Renal failure (N17-N19) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 18.0 | 14.2 | * | * | * | 1.9 | 6.3 | 20.3 | 46.2 | 98.7 | 222.6 | 443.8 |
| 1999..... | 17.8 | 8.6 | * | * | * | 1.8 | 6.3 | 19.4 | 41.5 | 97.5 | 239.7 | 476.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 20.3 | 9.1 | * | * | * | 1.7 | 4.1 | 11.6 | 36.3 | 94.3 | 179.8 | 383.7 |
| 1999..... | 19.9 | 11.8 | * | * | * | 2.0 | 4.4 | 11.9 | 33.6 | 88.3 | 169.0 | 405.5 |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS RACE, SEX, AND YEAR | 5-14 YEARS RACE, SEX, AND YEAR | 15-24 YEARS RACE, SEX, AND YEAR | 25-34 YEARS RACE, SEX, AND YEAR | 35-44 YEARS RACE, SEX, AND YEAR | 45-54 YEARS RACE, SEX, AND YEAR | 55-64 YEARS RACE, SEX, AND YEAR | 65-74 YEARS RACE, SEX, AND YEAR | 75-84 YEARS RACE, SEX, AND YEAR | 85 YEARS AND OVER RACE, SEX, AND YEAR |
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|
|---------------------------------------|--|-------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|---|

ALL RACES 3/, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALEWHITE, BOTH SEXES

WHITE, MALE

BLACK, BOTH SEXES

BLACK, MALE

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Pregnancy, childbirth and the puerperium (000-099) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.5 | 0.3 | * | ... | ... | ... | ... |
| 1999..... | 0.1 | ... | ... | * | 0.3 | 0.5 | 0.3 | * | ... | ... | ... | ... |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | ... | ... | * | 0.5 | 1.0 | 0.5 | * | ... | ... | ... | ... |
| 1999..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | ... | ... | ... | ... |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | ... | ... | ... | ... |
| 1999..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.2 | * | ... | ... | ... | ... |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.4 | * | ... | ... | ... | ... |
| 1999..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.4 | * | ... | ... | ... | ... |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | ... | ... | * | 0.6 | 1.2 | 0.7 | * | ... | ... | ... | ... |
| 1999..... | 0.5 | ... | ... | * | 0.8 | 1.3 | 0.7 | * | ... | ... | ... | ... |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.8 | ... | ... | * | 1.2 | 2.4 | 1.3 | * | ... | ... | ... | ... |
| 1999..... | 0.9 | ... | ... | * | 1.6 | 2.6 | 1.4 | * | ... | ... | ... | ... |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Pregnancy with abortive outcome (000-007) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.0 | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| 1999..... | 0.0 | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.0 | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| 1999..... | 0.0 | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| 1999..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| 1999..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| 1999..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |
| 1999..... | * | ... | ... | * | * | * | * | * | ... | ... | ... | ... |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other complications of pregnancy, childbirth and the puerperium (010-099) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | ... | ... | ... | ... |
| 1999..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.3 | * | ... | ... | ... | ... |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | ... | ... | ... | ... |
| 1999..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | ... | ... | ... | ... |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | ... | ... | ... | ... |
| 1999..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.2 | * | ... | ... | ... | ... |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.4 | * | ... | ... | ... | ... |
| 1999..... | 0.2 | ... | ... | * | 0.3 | 0.6 | 0.4 | * | ... | ... | ... | ... |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | ... | ... | * | 0.5 | 1.1 | 0.6 | * | ... | ... | ... | ... |
| 1999..... | 0.4 | ... | ... | * | 0.7 | 1.2 | 0.7 | * | ... | ... | ... | ... |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.7 | ... | ... | * | 1.1 | 2.0 | 1.1 | * | ... | ... | ... | ... |
| 1999..... | 0.8 | ... | ... | * | 1.5 | 2.3 | 1.3 | * | ... | ... | ... | ... |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.8 | 149.3 | 3.3 | 1.0 | 1.1 | 1.3 | 1.3 | 1.6 | 2.4 | 2.6 | 4.0 | 7.4 |
| 1999..... | 3.8 | 143.3 | 3.6 | 1.1 | 1.1 | 1.2 | 1.2 | 1.8 | 2.4 | 2.6 | 4.0 | 7.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 4.1 | 154.1 | 3.3 | 1.0 | 1.4 | 1.5 | 1.4 | 1.9 | 2.3 | 2.7 | 4.0 | 10.1 |
| 1999..... | 4.0 | 147.3 | 4.0 | 1.1 | 1.4 | 1.3 | 1.3 | 2.0 | 2.5 | 2.5 | 4.2 | 8.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.6 | 144.2 | 3.2 | 1.0 | 0.9 | 1.1 | 1.1 | 1.4 | 2.4 | 2.4 | 4.0 | 6.3 |
| 1999..... | 3.6 | 139.1 | 3.2 | 1.1 | 0.9 | 1.2 | 1.1 | 1.6 | 2.2 | 2.6 | 3.9 | 7.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.7 | 144.2 | 2.9 | 0.9 | 1.1 | 1.3 | 1.2 | 1.7 | 2.4 | 2.6 | 4.1 | 7.8 |
| 1999..... | 3.7 | 139.1 | 3.2 | 1.0 | 1.1 | 1.3 | 1.2 | 1.8 | 2.5 | 2.6 | 4.1 | 8.0 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 3.9 | 149.3 | 2.9 | 0.9 | 1.3 | 1.5 | 1.4 | 1.9 | 2.4 | 2.6 | 4.2 | 10.6 |
| 1999..... | 3.9 | 144.6 | 3.5 | 0.9 | 1.3 | 1.3 | 1.3 | 2.0 | 2.6 | 2.6 | 4.1 | 8.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 3.5 | 138.7 | 2.8 | 1.0 | 0.9 | 1.1 | 1.1 | 1.5 | 2.4 | 2.6 | 4.1 | 6.5 |
| 1999..... | 3.5 | 133.4 | 2.8 | 1.0 | 0.9 | 1.2 | 1.2 | 1.7 | 2.3 | 2.6 | 4.1 | 7.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.1 | 189.2 | 5.7 | 1.5 | 1.6 | 1.5 | 1.6 | 1.6 | 2.4 | 2.5 | 2.8 | * |
| 1999..... | 4.9 | 180.2 | 6.2 | 1.7 | 1.4 | 1.2 | 1.4 | 1.9 | 2.0 | 2.7 | 4.3 | 7.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|-------|-----|-----|-----|-----|-----|-----|-----|-----|---|---|
| 2000..... | 5.6 | 191.3 | 6.2 | 1.6 | 2.0 | 1.9 | 1.9 | 2.2 | * | 3.6 | * | * |
| 1999..... | 5.4 | 183.7 | 7.0 | 1.7 | 1.8 | 1.3 | 1.6 | 2.5 | 2.2 | * | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-------|-----|-----|-----|-----|-----|-----|-----|-----|---|---|
| 2000..... | 4.6 | 187.0 | 5.1 | 1.3 | 1.1 | 1.1 | 1.4 | 1.2 | 3.0 | * | * | * |
| 1999..... | 4.5 | 176.6 | 5.4 | 1.7 | 1.1 | 1.0 | 1.2 | 1.5 | 1.8 | 3.1 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

ALL AGES 1/
UNDER 1 YEAR 2/
1-4 YEARS
5-14 YEARS
15-24 YEARS
25-34 YEARS
35-44 YEARS
45-54 YEARS
55-64 YEARS
65-74 YEARS
75-84 YEARS
85 YEARS AND OVER

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 11.6 | 94.6 | 1.6 | 0.4 | 2.3 | 4.2 | 6.7 | 8.5 | 9.4 | 16.7 | 40.8 | 205.4 |
| 1999..... | 9.8 | 92.6 | 1.6 | 0.3 | 1.6 | 2.9 | 4.6 | 6.0 | 8.0 | 14.8 | 36.3 | 186.8 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|------|-------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| 2000..... | 11.8 | 109.7 | 1.8 | 0.5 | 3.0 | 5.7 | 8.9 | 11.7 | 12.5 | 20.8 | 45.2 | 173.4 |
| 1999..... | 9.7 | 106.1 | 1.8 | 0.4 | 2.2 | 3.8 | 6.0 | 8.3 | 10.6 | 18.5 | 39.6 | 160.0 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 11.4 | 78.8 | 1.5 | 0.4 | 1.5 | 2.6 | 4.4 | 5.4 | 6.6 | 13.4 | 37.9 | 219.2 |
| 1999..... | 9.9 | 78.5 | 1.4 | 0.3 | 1.0 | 2.0 | 3.2 | 3.9 | 5.5 | 11.8 | 34.1 | 198.1 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 11.6 | 75.7 | 1.5 | 0.4 | 2.2 | 4.0 | 6.5 | 7.8 | 8.7 | 15.9 | 40.3 | 209.9 |
| 1999..... | 9.6 | 73.6 | 1.3 | 0.3 | 1.4 | 2.6 | 4.1 | 5.2 | 7.2 | 14.1 | 35.3 | 189.9 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| 2000..... | 11.5 | 87.9 | 1.7 | 0.4 | 2.9 | 5.5 | 8.9 | 10.8 | 11.3 | 19.5 | 44.7 | 178.2 |
| 1999..... | 9.1 | 85.3 | 1.6 | 0.3 | 2.0 | 3.5 | 5.4 | 7.0 | 9.5 | 17.5 | 38.3 | 162.1 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 11.7 | 62.9 | 1.3 | 0.3 | 1.3 | 2.4 | 4.1 | 4.9 | 6.2 | 12.8 | 37.3 | 223.4 |
| 1999..... | 10.1 | 61.3 | 1.1 | 0.3 | 0.9 | 1.8 | 2.7 | 3.4 | 5.1 | 11.3 | 33.3 | 201.5 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|------|-------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| 2000..... | 14.2 | 208.1 | 2.6 | 0.7 | 3.2 | 6.0 | 9.3 | 15.1 | 17.6 | 28.2 | 56.5 | 186.9 |
| 1999..... | 13.2 | 205.2 | 3.1 | 0.5 | 2.7 | 5.0 | 9.1 | 13.6 | 16.1 | 24.0 | 55.1 | 182.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---------------------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|-------|-----|-----|-----|-----|------|------|------|------|------|-------|
| 2000..... | 16.1 | 239.5 | 2.6 | 0.9 | 3.7 | 8.0 | 10.9 | 21.5 | 26.3 | 38.2 | 63.2 | 149.0 |
| 1999..... | 15.2 | 231.4 | 3.1 | 0.6 | 3.5 | 7.0 | 11.6 | 20.4 | 24.1 | 31.2 | 64.5 | 170.6 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|-------|-----|---|-----|-----|-----|-----|------|------|------|-------|
| 2000..... | 12.5 | 175.5 | 2.6 | * | 2.7 | 4.3 | 7.8 | 9.8 | 10.9 | 21.0 | 52.5 | 202.8 |
| 1999..... | 11.3 | 178.1 | 3.0 | * | 1.9 | 3.2 | 6.9 | 8.0 | 10.0 | 18.7 | 49.5 | 187.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 64.8 | 28.7 | 4.0 | 2.2 | 4.6 | 8.3 | 18.1 | 35.9 | 63.2 | 137.6 | 390.0 | 1,426.8 |
| 1999..... | 63.0 | 30.1 | 4.3 | 2.3 | 4.7 | 8.1 | 17.8 | 34.6 | 61.5 | 138.1 | 375.4 | 1,403.5 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 53.7 | 32.0 | 4.4 | 2.3 | 5.4 | 9.7 | 21.5 | 43.5 | 72.0 | 150.2 | 384.9 | 1,201.4 |
| 1999..... | 52.4 | 34.8 | 4.6 | 2.4 | 5.4 | 9.5 | 21.7 | 41.6 | 70.6 | 150.4 | 379.7 | 1,179.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 75.5 | 25.3 | 3.7 | 2.1 | 3.8 | 6.8 | 14.7 | 28.5 | 55.2 | 127.3 | 393.4 | 1,523.9 |
| 1999..... | 73.1 | 25.1 | 4.0 | 2.3 | 3.9 | 6.7 | 14.0 | 27.9 | 53.3 | 128.1 | 372.5 | 1,498.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 68.1 | 24.4 | 3.5 | 2.1 | 4.3 | 7.4 | 16.0 | 32.3 | 59.1 | 134.6 | 387.9 | 1,448.3 |
| 1999..... | 66.2 | 24.8 | 4.0 | 2.2 | 4.4 | 7.4 | 15.6 | 30.8 | 57.7 | 134.3 | 373.8 | 1,431.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 55.2 | 26.4 | 3.9 | 2.3 | 5.2 | 9.0 | 19.5 | 39.8 | 66.8 | 145.6 | 380.7 | 1,215.9 |
| 1999..... | 53.8 | 28.8 | 4.3 | 2.2 | 5.2 | 9.1 | 19.7 | 37.5 | 66.0 | 145.8 | 375.3 | 1,193.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 80.6 | 22.3 | 3.1 | 1.9 | 3.4 | 5.7 | 12.5 | 24.9 | 52.0 | 125.3 | 392.8 | 1,547.7 |
| 1999..... | 78.1 | 20.6 | 3.7 | 2.2 | 3.5 | 5.7 | 11.4 | 24.3 | 50.0 | 124.8 | 372.8 | 1,530.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 60.3 | 52.4 | 6.7 | 3.0 | 6.6 | 15.6 | 34.5 | 71.0 | 108.8 | 187.1 | 476.4 | 1,392.5 |
| 1999..... | 58.3 | 60.3 | 5.9 | 3.2 | 6.8 | 13.4 | 35.7 | 70.0 | 105.1 | 195.7 | 449.5 | 1,288.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|------|------|------|-------|-------|-------|---------|
| 2000..... | 56.4 | 63.1 | 6.9 | 2.7 | 7.0 | 16.3 | 39.0 | 84.4 | 136.1 | 218.3 | 508.4 | 1,222.8 |
| 1999..... | 54.8 | 68.8 | 6.0 | 3.8 | 7.1 | 13.9 | 39.6 | 84.3 | 127.2 | 217.4 | 493.0 | 1,197.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|------|------|------|------|-------|-------|---------|
| 2000..... | 63.8 | 41.2 | 6.4 | 3.3 | 6.2 | 15.0 | 30.5 | 59.7 | 88.1 | 164.4 | 457.0 | 1,463.4 |
| 1999..... | 61.5 | 51.5 | 5.7 | 2.7 | 6.4 | 12.9 | 32.2 | 58.1 | 88.2 | 179.9 | 423.4 | 1,326.9 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Accidents (unintentional injuries) (V01-X59,Y85-Y86) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 35.6 | 22.9 | 12.1 | 7.5 | 36.8 | 31.4 | 34.3 | 33.0 | 31.3 | 42.4 | 95.5 | 269.9 |
| 1999..... | 35.9 | 22.1 | 12.6 | 7.8 | 36.2 | 31.3 | 34.0 | 32.5 | 31.1 | 45.1 | 101.1 | 280.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 47.4 | 25.5 | 14.2 | 9.2 | 53.3 | 48.7 | 51.3 | 49.9 | 44.9 | 56.5 | 122.2 | 341.2 |
| 1999..... | 47.7 | 25.3 | 14.6 | 9.8 | 51.9 | 48.3 | 50.5 | 49.2 | 45.0 | 60.9 | 130.8 | 361.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 24.2 | 20.2 | 9.8 | 5.8 | 19.5 | 14.4 | 17.6 | 16.9 | 18.9 | 30.7 | 77.5 | 239.2 |
| 1999..... | 24.6 | 18.8 | 10.4 | 5.8 | 19.6 | 14.6 | 17.7 | 16.6 | 18.5 | 32.1 | 81.2 | 247.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 36.5 | 18.8 | 11.1 | 7.1 | 38.8 | 31.9 | 34.3 | 31.6 | 30.4 | 41.7 | 96.7 | 279.6 |
| 1999..... | 36.6 | 18.6 | 11.4 | 7.3 | 37.9 | 31.7 | 33.8 | 31.0 | 29.6 | 44.0 | 101.5 | 289.9 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 48.0 | 21.3 | 13.1 | 8.6 | 56.0 | 49.2 | 51.1 | 47.3 | 42.7 | 55.0 | 122.2 | 352.7 |
| 1999..... | 47.9 | 21.5 | 13.2 | 9.0 | 53.9 | 48.7 | 50.1 | 46.3 | 42.2 | 58.9 | 130.4 | 371.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 25.4 | 16.1 | 9.0 | 5.5 | 20.6 | 14.4 | 17.4 | 16.3 | 19.1 | 30.6 | 79.4 | 248.4 |
| 1999..... | 25.7 | 15.5 | 9.6 | 5.5 | 20.8 | 14.6 | 17.5 | 16.0 | 17.9 | 31.7 | 82.0 | 255.6 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 34.8 | 47.0 | 18.2 | 10.1 | 29.0 | 33.8 | 40.1 | 48.1 | 41.3 | 51.1 | 86.4 | 181.6 |
| 1999..... | 36.5 | 44.3 | 19.8 | 11.3 | 29.9 | 33.4 | 40.4 | 49.4 | 45.5 | 56.7 | 105.3 | 208.0 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|------|------|------|------|------|------|------|------|------|-------|-------|
| 2000..... | 50.9 | 50.3 | 21.1 | 13.0 | 43.0 | 54.0 | 62.5 | 78.5 | 71.9 | 75.9 | 128.8 | 254.3 |
| 1999..... | 53.0 | 48.8 | 23.5 | 14.9 | 45.1 | 53.3 | 62.6 | 82.5 | 74.4 | 84.8 | 146.4 | 293.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|------|-----|------|------|------|------|------|------|------|-------|
| 2000..... | 20.2 | 43.7 | 15.2 | 7.0 | 14.9 | 15.6 | 20.1 | 22.8 | 18.1 | 33.1 | 60.7 | 151.2 |
| 1999..... | 21.6 | 39.7 | 15.9 | 7.7 | 14.6 | 15.5 | 20.8 | 21.8 | 23.4 | 36.1 | 80.6 | 172.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Transport accidents (V01-V99,Y85) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 17.0 | 4.5 | 4.6 | 4.8 | 28.7 | 19.8 | 17.1 | 16.1 | 16.2 | 18.2 | 27.1 | 31.1 |
| 1999..... | 17.0 | 4.9 | 4.7 | 5.0 | 28.3 | 19.6 | 16.9 | 15.8 | 16.3 | 19.6 | 28.6 | 31.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 23.9 | 4.8 | 5.1 | 5.6 | 40.2 | 29.9 | 24.9 | 23.4 | 22.9 | 24.6 | 37.4 | 56.8 |
| 1999..... | 23.7 | 5.2 | 5.2 | 6.0 | 39.1 | 29.3 | 24.2 | 22.9 | 23.0 | 26.4 | 40.2 | 61.1 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.4 | 4.2 | 4.1 | 4.0 | 16.8 | 9.7 | 9.3 | 9.0 | 10.1 | 13.0 | 20.2 | 20.0 |
| 1999..... | 10.6 | 4.7 | 4.2 | 4.0 | 17.0 | 10.1 | 9.7 | 8.9 | 10.2 | 14.1 | 20.7 | 19.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 17.3 | 4.0 | 4.3 | 4.9 | 30.5 | 19.7 | 17.1 | 15.7 | 16.1 | 18.1 | 27.6 | 31.9 |
| 1999..... | 17.3 | 4.5 | 4.1 | 5.0 | 29.8 | 19.8 | 16.8 | 15.4 | 16.0 | 19.6 | 28.8 | 32.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 24.1 | 4.4 | 4.7 | 5.6 | 42.3 | 29.6 | 24.8 | 22.9 | 22.3 | 24.1 | 37.5 | 58.8 |
| 1999..... | 23.9 | 5.2 | 4.4 | 5.9 | 40.9 | 29.3 | 23.9 | 22.3 | 22.3 | 26.2 | 40.2 | 61.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.7 | 3.6 | 3.8 | 4.1 | 17.9 | 9.7 | 9.4 | 8.7 | 10.4 | 13.1 | 20.8 | 20.3 |
| 1999..... | 10.8 | 3.9 | 3.8 | 4.0 | 18.1 | 10.2 | 9.6 | 8.7 | 10.1 | 14.2 | 21.1 | 19.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.5 | 6.9 | 6.3 | 5.1 | 22.0 | 22.3 | 18.7 | 19.8 | 17.6 | 19.1 | 20.3 | 21.6 |
| 1999..... | 16.9 | 7.7 | 7.9 | 5.8 | 22.4 | 20.6 | 19.2 | 19.7 | 18.6 | 18.9 | 26.5 | 27.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Transport accidents (V01-V99,Y85) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 25.0 | 7.1 | 7.0 | 6.2 | 31.7 | 35.8 | 30.0 | 30.8 | 30.8 | 30.8 | 35.3 | 41.5 |
| 1999..... | 25.1 | * | 8.9 | 7.3 | 32.1 | 33.1 | 29.3 | 31.8 | 30.3 | 28.0 | 42.3 | 63.0 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 8.9 | * | 5.7 | 3.9 | 12.2 | 10.1 | 8.8 | 10.6 | 7.5 | 10.7 | 11.3 | 13.3 |
| 1999..... | 9.5 | 9.3 | 6.9 | 4.3 | 12.5 | 9.4 | 10.2 | 9.6 | 9.7 | 12.4 | 17.1 | 13.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 15.7 | 4.4 | 4.3 | 4.5 | 27.5 | 18.4 | 15.4 | 14.4 | 14.6 | 16.7 | 25.8 | 30.0 |
| 1999..... | 15.5 | 4.8 | 4.3 | 4.5 | 26.8 | 17.9 | 15.0 | 13.9 | 14.4 | 18.0 | 26.7 | 30.2 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 21.9 | 4.6 | 4.8 | 5.1 | 38.3 | 27.6 | 22.1 | 20.8 | 20.2 | 22.0 | 35.0 | 54.6 |
| 1999..... | 21.4 | 5.0 | 4.7 | 5.3 | 36.9 | 26.5 | 21.3 | 19.8 | 19.7 | 23.6 | 37.5 | 57.6 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 9.9 | 4.1 | 3.8 | 3.8 | 16.3 | 9.3 | 8.8 | 8.3 | 9.6 | 12.4 | 19.5 | 19.4 |
| 1999..... | 9.9 | 4.7 | 3.8 | 3.7 | 16.3 | 9.3 | 8.9 | 8.2 | 9.6 | 13.4 | 19.5 | 18.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.0 | 3.8 | 4.0 | 4.5 | 29.2 | 18.2 | 15.4 | 14.1 | 14.5 | 16.6 | 26.2 | 30.7 |
| 1999..... | 15.7 | 4.4 | 3.8 | 4.4 | 28.3 | 18.0 | 14.9 | 13.5 | 14.1 | 18.0 | 27.0 | 30.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 22.0 | 4.1 | 4.5 | 5.1 | 40.3 | 27.1 | 21.9 | 20.2 | 19.5 | 21.5 | 35.0 | 56.4 |
| 1999..... | 21.5 | 5.0 | 4.1 | 5.2 | 38.6 | 26.4 | 20.9 | 19.2 | 19.0 | 23.4 | 37.5 | 58.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 10.2 | 3.4 | 3.6 | 3.8 | 17.3 | 9.3 | 8.9 | 8.1 | 9.8 | 12.5 | 20.1 | 19.8 |
| 1999..... | 10.1 | 3.8 | 3.4 | 3.7 | 17.3 | 9.4 | 8.8 | 8.0 | 9.4 | 13.5 | 19.9 | 19.0 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 15.6 | 6.9 | 6.0 | 4.8 | 21.1 | 21.2 | 17.3 | 18.3 | 16.3 | 17.7 | 19.2 | 20.4 |
| 1999..... | 15.7 | 7.6 | 7.2 | 5.4 | 21.4 | 19.3 | 17.6 | 17.7 | 16.8 | 17.6 | 24.5 | 25.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|-----|-----|-----|------|------|------|------|------|------|------|------|
| 2000..... | 23.4 | 7.1 | 6.5 | 5.8 | 30.2 | 33.9 | 27.4 | 28.2 | 28.3 | 28.1 | 33.5 | 40.4 |
| 1999..... | 23.1 | * | 7.9 | 6.5 | 30.7 | 30.8 | 26.5 | 28.4 | 26.7 | 25.4 | 40.2 | 56.5 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|------|-----|-----|------|-----|------|------|------|
| 2000..... | 8.5 | * | 5.5 | 3.8 | 12.0 | 9.8 | 8.3 | 10.1 | 7.2 | 10.2 | 10.5 | 12.0 |
| 1999..... | 9.0 | 9.3 | 6.5 | 4.2 | 12.1 | 8.9 | 9.7 | 8.9 | 9.2 | 11.8 | 15.1 | 12.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.5 | * | 0.2 | 0.2 | 0.6 | 0.6 | 0.7 | 0.6 | 0.5 | 0.6 | 0.7 | 0.8 |
| 1999..... | 0.8 | * | 0.3 | 0.3 | 1.0 | 0.8 | 0.9 | 0.8 | 0.8 | 0.8 | 1.3 | 1.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.8 | * | * | 0.3 | 1.0 | 0.9 | 1.2 | 0.9 | 0.9 | 0.9 | 1.1 | * |
| 1999..... | 1.2 | * | 0.3 | 0.5 | 1.4 | 1.3 | 1.4 | 1.3 | 1.2 | 1.2 | 1.8 | 2.8 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 0.2 | 0.3 | 0.5 | * |
| 1999..... | 0.4 | * | 0.3 | 0.2 | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 1.0 | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.5 | * | 0.2 | 0.3 | 0.7 | 0.6 | 0.7 | 0.6 | 0.5 | 0.5 | 0.7 | 0.8 |
| 1999..... | 0.8 | * | 0.3 | 0.3 | 1.0 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 1.3 | 1.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.8 | * | * | 0.3 | 1.0 | 0.9 | 1.1 | 0.9 | 0.8 | 0.9 | 1.2 | * |
| 1999..... | 1.2 | * | * | 0.4 | 1.4 | 1.3 | 1.4 | 1.3 | 1.2 | 1.2 | 1.8 | 2.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | * | 0.3 | 0.4 | * |
| 1999..... | 0.4 | * | * | 0.2 | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.9 | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.6 | * | * | * | 0.5 | 0.6 | 0.9 | 0.8 | * | * | * | * |
| 1999..... | 0.7 | * | * | 0.3 | 0.6 | 0.8 | 0.9 | 1.1 | * | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 0.7 | * | * | 0.1 | 0.6 | 0.8 | 0.9 | 1.0 | 1.1 | 1.0 | 0.6 | * |
| 1999..... | 0.7 | * | * | 0.2 | 0.5 | 0.9 | 0.9 | 1.0 | 1.1 | 0.9 | 0.5 | * |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 1.2 | * | * | 0.2 | 0.9 | 1.4 | 1.6 | 1.7 | 1.9 | 1.8 | 1.2 | * |
| 1999..... | 1.1 | * | * | 0.3 | 0.8 | 1.5 | 1.4 | 1.8 | 2.0 | 1.6 | 0.9 | * |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 | * | * |
| 1999..... | 0.3 | * | * | 0.1 | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | * |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 0.8 | * | * | 0.2 | 0.6 | 0.9 | 1.0 | 1.0 | 1.1 | 1.0 | 0.7 | * |
| 1999..... | 0.7 | * | * | 0.2 | 0.6 | 1.0 | 1.0 | 1.1 | 1.1 | 0.9 | 0.5 | * |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 1.3 | * | * | 0.2 | 1.0 | 1.6 | 1.8 | 1.8 | 2.0 | 1.8 | 1.3 | * |
| 1999..... | 1.2 | * | * | 0.3 | 0.9 | 1.6 | 1.5 | 1.8 | 2.0 | 1.7 | 0.9 | * |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|-----|-----|---|---|
| 2000..... | 0.2 | * | * | * | 0.3 | 0.2 | 0.3 | 0.3 | 0.4 | 0.3 | * | * |
| 1999..... | 0.3 | * | * | 0.1 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | * | * |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|---|---|---|
| 2000..... | 0.4 | * | * | * | * | 0.5 | 0.5 | 0.6 | * | * | * | * |
| 1999..... | 0.5 | * | * | * | 0.3 | 0.5 | 0.7 | 0.9 | 1.1 | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------------|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Nontransport accidents (W00-X59,Y86) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 18.6 | 18.4 | 7.5 | 2.7 | 8.0 | 11.7 | 17.3 | 17.0 | 15.1 | 24.1 | 68.3 | 238.8 |
| 1999..... | 18.9 | 17.2 | 7.8 | 2.8 | 7.9 | 11.7 | 17.1 | 16.7 | 14.8 | 25.4 | 72.6 | 249.1 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 23.5 | 20.7 | 9.1 | 3.6 | 13.1 | 18.8 | 26.4 | 26.4 | 22.0 | 31.9 | 84.7 | 284.5 |
| 1999..... | 23.9 | 20.1 | 9.4 | 3.8 | 12.8 | 19.0 | 26.4 | 26.2 | 22.0 | 34.5 | 90.6 | 300.1 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 13.9 | 15.9 | 5.8 | 1.7 | 2.7 | 4.7 | 8.3 | 8.0 | 8.8 | 17.7 | 57.3 | 219.2 |
| 1999..... | 14.0 | 14.1 | 6.2 | 1.8 | 2.6 | 4.6 | 8.0 | 7.7 | 8.3 | 18.0 | 60.5 | 227.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 19.2 | 14.8 | 6.8 | 2.2 | 8.4 | 12.2 | 17.2 | 15.9 | 14.3 | 23.6 | 69.2 | 247.8 |
| 1999..... | 19.4 | 14.0 | 7.3 | 2.3 | 8.0 | 11.9 | 17.1 | 15.5 | 13.6 | 24.4 | 72.7 | 257.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 23.8 | 17.0 | 8.4 | 3.0 | 13.6 | 19.5 | 26.3 | 24.4 | 20.4 | 30.9 | 84.7 | 294.0 |
| 1999..... | 24.0 | 16.3 | 8.7 | 3.1 | 13.0 | 19.4 | 26.2 | 24.0 | 20.0 | 32.7 | 90.3 | 309.9 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 14.8 | 12.5 | 5.2 | 1.4 | 2.7 | 4.7 | 8.0 | 7.6 | 8.7 | 17.4 | 58.6 | 228.0 |
| 1999..... | 14.8 | 11.6 | 5.8 | 1.5 | 2.7 | 4.4 | 7.8 | 7.3 | 7.8 | 17.5 | 60.9 | 235.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 18.2 | 40.2 | 11.9 | 5.0 | 7.1 | 11.5 | 21.3 | 28.3 | 23.8 | 32.0 | 66.1 | 159.9 |
| 1999..... | 19.6 | 36.6 | 11.9 | 5.5 | 7.6 | 12.8 | 21.2 | 29.7 | 26.9 | 37.7 | 78.7 | 180.2 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Nontransport accidents (W00-X59,Y86)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|------|------|-----|------|------|------|------|------|------|-------|-------|
| 2000..... | 25.8 | 43.2 | 14.2 | 6.8 | 11.4 | 18.1 | 32.6 | 47.7 | 41.1 | 45.2 | 93.5 | 212.8 |
| 1999..... | 27.8 | 42.5 | 14.6 | 7.6 | 13.0 | 20.2 | 33.2 | 50.7 | 44.2 | 56.8 | 104.1 | 230.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|-----|-----|------|------|------|------|------|-------|
| 2000..... | 11.4 | 37.1 | 9.5 | 3.2 | 2.7 | 5.5 | 11.4 | 12.2 | 10.5 | 22.4 | 49.5 | 137.8 |
| 1999..... | 12.1 | 30.4 | 9.0 | 3.4 | 2.1 | 6.1 | 10.6 | 12.3 | 13.7 | 23.8 | 63.5 | 159.3 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Falls (W00-W19) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.8 | * | 0.2 | 0.1 | 0.6 | 0.8 | 1.4 | 2.3 | 4.0 | 9.1 | 31.2 | 111.1 |
| 1999..... | 4.8 | * | 0.4 | 0.1 | 0.6 | 0.9 | 1.4 | 2.3 | 3.8 | 9.1 | 31.7 | 109.7 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 5.3 | * | 0.3 | 0.1 | 1.0 | 1.5 | 2.2 | 3.8 | 5.9 | 12.1 | 39.2 | 138.0 |
| 1999..... | 5.3 | * | 0.5 | 0.2 | 1.1 | 1.6 | 2.4 | 3.8 | 5.8 | 12.5 | 40.0 | 137.2 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.4 | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 2.1 | 6.7 | 25.8 | 99.5 |
| 1999..... | 4.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 1.9 | 6.3 | 26.2 | 98.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.4 | * | 0.2 | 0.1 | 0.7 | 0.9 | 1.4 | 2.3 | 3.9 | 9.3 | 32.7 | 117.8 |
| 1999..... | 5.3 | * | 0.4 | 0.1 | 0.7 | 0.9 | 1.5 | 2.2 | 3.6 | 9.2 | 33.1 | 115.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 5.8 | * | * | 0.2 | 1.2 | 1.6 | 2.2 | 3.7 | 5.8 | 12.3 | 40.7 | 145.7 |
| 1999..... | 5.8 | * | 0.5 | 0.2 | 1.2 | 1.6 | 2.4 | 3.6 | 5.5 | 12.5 | 41.4 | 144.5 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 5.0 | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 2.1 | 6.8 | 27.2 | 105.8 |
| 1999..... | 4.9 | * | * | * | 0.2 | 0.3 | 0.6 | 0.8 | 1.8 | 6.5 | 27.5 | 103.7 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 1.3 | 2.8 | 4.8 | 8.1 | 15.2 | 38.9 |
| 1999..... | 2.5 | * | * | * | 0.4 | 0.8 | 1.3 | 3.5 | 5.6 | 9.0 | 17.4 | 45.4 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|-----------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Falls (W00-W19) | | | | | | | | | | | | |
| BLACK, MALE | | | | | | | | | | | | |
| 2000..... | 2.9 | * | * | * | * | * | 2.1 | 4.8 | 8.5 | 10.9 | 20.6 | 56.4 |
| 1999..... | 3.4 | * | * | * | * | 1.4 | 2.5 | 6.1 | 9.2 | 14.1 | 23.7 | 54.3 |
| BLACK, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.5 | * | * | * | * | * | * | 1.0 | 2.0 | 6.0 | 12.0 | 31.6 |
| 1999..... | 1.7 | * | * | * | * | * | * | 1.4 | 2.9 | 5.3 | 13.7 | 41.6 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Accidental discharge of firearms (W32-W34) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | 0.2 | 0.5 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | * |
| 1999..... | 0.3 | * | * | 0.2 | 0.7 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | 0.3 | 0.9 | 0.6 | 0.6 | 0.4 | 0.4 | 0.4 | 0.5 | * |
| 1999..... | 0.5 | * | * | 0.3 | 1.2 | 0.6 | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | * |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | * | 0.1 | 0.1 | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | 0.1 | 0.1 | 0.1 | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | 0.2 | 0.5 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | * |
| 1999..... | 0.3 | * | * | 0.2 | 0.6 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | * | 0.3 | 0.8 | 0.5 | 0.6 | 0.4 | 0.4 | 0.4 | 0.5 | * |
| 1999..... | 0.5 | * | * | 0.3 | 1.0 | 0.6 | 0.5 | 0.5 | 0.4 | 0.4 | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.4 | * | * | * | 0.9 | 0.6 | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | 1.2 | 0.4 | * | * | * | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Accidental drowning and submersion (W65-W74) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | 1.9 | 3.3 | 0.9 | 1.7 | 1.1 | 1.1 | 1.0 | 0.9 | 1.0 | 1.3 | 1.5 |
| 1999..... | 1.3 | 1.8 | 3.2 | 0.9 | 1.7 | 1.2 | 1.1 | 1.0 | 0.9 | 1.0 | 1.3 | 1.6 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 2.0 | 2.2 | 4.1 | 1.3 | 3.0 | 2.0 | 1.8 | 1.6 | 1.6 | 1.6 | 1.8 | 2.6 |
| 1999..... | 2.1 | 2.0 | 4.0 | 1.4 | 3.1 | 2.1 | 1.8 | 1.8 | 1.5 | 1.6 | 2.3 | 2.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.5 | 1.7 | 2.4 | 0.6 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.5 | 0.9 | 1.0 |
| 1999..... | 0.5 | 1.6 | 2.4 | 0.5 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | 0.6 | 0.6 | 1.4 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.2 | 1.6 | 3.5 | 0.7 | 1.5 | 1.1 | 1.0 | 0.9 | 0.9 | 1.0 | 1.2 | 1.4 |
| 1999..... | 1.2 | 1.5 | 3.5 | 0.6 | 1.5 | 1.1 | 1.1 | 1.0 | 0.9 | 1.0 | 1.3 | 1.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.9 | 1.8 | 4.4 | 0.9 | 2.6 | 1.9 | 1.7 | 1.5 | 1.6 | 1.6 | 1.8 | 2.6 |
| 1999..... | 1.9 | 1.7 | 4.3 | 0.8 | 2.7 | 1.9 | 1.8 | 1.8 | 1.4 | 1.5 | 2.2 | 2.3 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.5 | * | 2.5 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.5 | 0.8 | 1.0 |
| 1999..... | 0.5 | 1.4 | 2.6 | 0.4 | 0.3 | 0.2 | 0.4 | 0.3 | 0.4 | 0.6 | 0.6 | 1.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.7 | 4.5 | 2.5 | 2.2 | 2.3 | 1.1 | 1.4 | 1.3 | * | * | * | * |
| 1999..... | 1.8 | 3.7 | 2.7 | 2.6 | 2.4 | 1.7 | 1.1 | 0.9 | 1.2 | 1.3 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Accidental exposure to smoke, fire and flames (X00-X09) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.2 | 1.0 | 1.9 | 0.7 | 0.5 | 0.6 | 0.9 | 1.2 | 1.5 | 2.2 | 3.8 | 6.1 |
| 1999..... | 1.2 | 1.1 | 2.0 | 0.7 | 0.5 | 0.7 | 0.9 | 1.0 | 1.4 | 2.3 | 4.0 | 5.9 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.5 | 1.0 | 2.2 | 0.7 | 0.6 | 0.8 | 1.2 | 1.7 | 2.1 | 2.8 | 4.9 | 8.9 |
| 1999..... | 1.5 | 1.1 | 2.2 | 0.7 | 0.7 | 1.0 | 1.3 | 1.5 | 1.7 | 3.0 | 5.1 | 8.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | 1.6 | 0.6 | 0.4 | 0.4 | 0.6 | 0.7 | 1.0 | 1.7 | 3.1 | 4.9 |
| 1999..... | 0.9 | 1.1 | 1.7 | 0.6 | 0.4 | 0.4 | 0.5 | 0.6 | 1.0 | 1.7 | 3.3 | 4.6 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.1 | 0.8 | 1.3 | 0.5 | 0.5 | 0.6 | 0.8 | 1.1 | 1.3 | 1.9 | 3.1 | 5.1 |
| 1999..... | 1.1 | 0.9 | 1.5 | 0.5 | 0.5 | 0.7 | 0.8 | 0.9 | 1.2 | 1.8 | 3.3 | 4.7 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.3 | * | 1.6 | 0.5 | 0.6 | 0.8 | 1.1 | 1.5 | 1.7 | 2.3 | 4.0 | 7.8 |
| 1999..... | 1.3 | * | 1.7 | 0.6 | 0.7 | 1.0 | 1.2 | 1.3 | 1.5 | 2.3 | 4.2 | 7.7 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.8 | * | 1.1 | 0.5 | 0.3 | 0.4 | 0.5 | 0.6 | 0.9 | 1.5 | 2.5 | 3.9 |
| 1999..... | 0.8 | * | 1.3 | 0.5 | 0.3 | 0.4 | 0.5 | 0.5 | 0.8 | 1.4 | 2.7 | 3.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.6 | * | 5.4 | 1.6 | 0.8 | 1.2 | 1.7 | 2.4 | 4.2 | 5.8 | 13.4 | 20.4 |
| 1999..... | 2.6 | * | 5.0 | 1.6 | 0.7 | 1.0 | 1.8 | 2.2 | 3.4 | 6.7 | 13.5 | 21.7 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Accidental exposure to smoke, fire and flames (X00-X09)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|-----|-----|-----|-----|-----|-----|-----|-----|------|------|
| 2000..... | 3.3 | * | 6.3 | 1.8 | 0.9 | 1.6 | 2.4 | 3.7 | 6.6 | 8.4 | 17.9 | 25.5 |
| 1999..... | 3.1 | * | 5.6 | 1.8 | 0.8 | 1.2 | 2.7 | 3.3 | 4.3 | 9.8 | 17.7 | 27.2 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|-----|-----|---|-----|-----|-----|-----|-----|------|------|
| 2000..... | 2.0 | * | 4.6 | 1.4 | * | 0.9 | 1.1 | 1.4 | 2.4 | 3.9 | 10.7 | 18.2 |
| 1999..... | 2.1 | * | 4.4 | 1.4 | * | 0.8 | 1.0 | 1.3 | 2.7 | 4.5 | 11.0 | 19.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Accidental poisoning and exposure to noxious substances (X40-X49) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | * | 0.2 | 0.1 | 3.0 | 6.4 | 10.4 | 8.2 | 2.9 | 1.5 | 2.0 | 4.3 |
| 1999..... | 4.5 | * | 0.2 | 0.1 | 2.6 | 6.2 | 10.2 | 7.9 | 2.9 | 1.6 | 2.2 | 3.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.8 | * | 0.3 | 0.1 | 4.7 | 9.8 | 15.2 | 12.2 | 3.5 | 1.8 | 2.1 | 5.9 |
| 1999..... | 6.7 | * | * | 0.1 | 3.9 | 9.7 | 15.0 | 12.0 | 4.0 | 2.1 | 2.4 | 4.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.6 | * | * | * | 1.3 | 3.0 | 5.7 | 4.4 | 2.3 | 1.3 | 1.9 | 3.6 |
| 1999..... | 2.4 | * | * | * | 1.1 | 2.8 | 5.3 | 4.0 | 1.8 | 1.3 | 2.0 | 3.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 4.6 | * | * | 0.1 | 3.4 | 6.8 | 10.4 | 7.5 | 2.6 | 1.5 | 2.1 | 4.0 |
| 1999..... | 4.4 | * | 0.2 | 0.1 | 2.8 | 6.5 | 10.0 | 7.2 | 2.5 | 1.5 | 2.2 | 3.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 6.8 | * | * | * | 5.4 | 10.5 | 15.3 | 11.0 | 2.9 | 1.6 | 2.1 | 5.6 |
| 1999..... | 6.6 | * | * | * | 4.3 | 10.2 | 15.0 | 10.7 | 3.3 | 1.9 | 2.4 | 4.4 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.6 | * | * | * | 1.3 | 3.1 | 5.5 | 4.2 | 2.4 | 1.4 | 2.0 | 3.4 |
| 1999..... | 2.3 | * | * | * | 1.2 | 2.7 | 5.1 | 3.8 | 1.8 | 1.2 | 2.1 | 2.8 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.7 | * | * | * | 1.4 | 5.8 | 13.1 | 15.9 | 5.6 | 2.5 | * | 8.2 |
| 1999..... | 5.9 | * | * | * | 1.6 | 6.4 | 13.3 | 16.0 | 6.1 | 3.0 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Accidental poisoning and exposure to noxious substances (X40-X49)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|------|------|------|-----|---|---|
| 2000..... | 8.5 | * | * | * | 1.7 | 8.7 | 18.9 | 25.9 | 10.0 | 4.3 | * | * |
| 1999..... | 9.0 | * | * | * | 2.7 | 9.5 | 19.6 | 27.2 | 11.5 | 4.2 | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|-----|---|---|
| 2000..... | 3.2 | * | * | * | 1.0 | 3.2 | 7.9 | 7.5 | 2.3 | * | * | * |
| 1999..... | 3.1 | * | * | * | * | 3.6 | 7.7 | 6.7 | 2.0 | 2.1 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 6.3 | 14.9 | 1.8 | 0.7 | 1.7 | 2.4 | 3.2 | 4.1 | 5.6 | 10.0 | 29.9 | 115.6 |
| 1999..... | 6.7 | 13.7 | 1.9 | 0.8 | 1.8 | 2.3 | 3.2 | 4.2 | 5.7 | 11.2 | 33.1 | 128.3 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 7.4 | 16.6 | 2.1 | 0.9 | 2.9 | 4.1 | 5.4 | 6.7 | 8.5 | 13.2 | 36.3 | 128.3 |
| 1999..... | 7.8 | 16.1 | 2.4 | 1.1 | 2.9 | 4.0 | 5.3 | 6.7 | 8.5 | 15.0 | 40.4 | 146.9 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|-----|------|-------|
| 2000..... | 5.3 | 13.0 | 1.4 | 0.4 | 0.5 | 0.7 | 1.1 | 1.5 | 3.0 | 7.5 | 25.5 | 110.1 |
| 1999..... | 5.8 | 11.2 | 1.5 | 0.5 | 0.5 | 0.7 | 1.2 | 1.7 | 3.1 | 8.0 | 28.3 | 120.4 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 6.7 | 11.9 | 1.6 | 0.7 | 1.8 | 2.5 | 3.2 | 3.9 | 5.4 | 9.7 | 29.9 | 119.2 |
| 1999..... | 7.1 | 11.1 | 1.8 | 0.8 | 1.9 | 2.4 | 3.3 | 3.9 | 5.3 | 10.7 | 32.7 | 132.1 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|------|------|-------|
| 2000..... | 7.6 | 13.7 | 2.0 | 1.0 | 3.0 | 4.3 | 5.4 | 6.3 | 8.0 | 12.6 | 35.6 | 131.3 |
| 1999..... | 8.0 | 12.7 | 2.1 | 1.1 | 3.1 | 4.1 | 5.4 | 6.3 | 7.9 | 14.2 | 39.7 | 150.4 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|-----|------|-------|
| 2000..... | 5.8 | 10.1 | 1.2 | 0.4 | 0.5 | 0.7 | 1.0 | 1.5 | 3.0 | 7.2 | 26.0 | 114.0 |
| 1999..... | 6.2 | 9.3 | 1.4 | 0.5 | 0.6 | 0.7 | 1.2 | 1.7 | 2.8 | 7.8 | 27.9 | 124.5 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|------|------|------|-------|
| 2000..... | 5.6 | 32.3 | 2.6 | 0.8 | 1.5 | 2.3 | 3.5 | 5.8 | 8.3 | 14.6 | 34.3 | 90.3 |
| 1999..... | 6.3 | 29.7 | 2.8 | 0.8 | 1.3 | 2.5 | 3.4 | 6.8 | 10.2 | 17.6 | 43.8 | 105.8 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| 2000..... | 7.6 | 33.7 | 2.8 | 1.1 | 2.5 | 3.8 | 6.0 | 10.5 | 14.5 | 19.4 | 51.5 | 114.9 |
| 1999..... | 8.3 | 35.3 | 3.8 | 1.0 | 2.4 | 4.1 | 5.8 | 11.6 | 16.5 | 25.9 | 55.8 | 142.4 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|---|---|-----|-----|-----|-----|------|------|------|
| 2000..... | 3.9 | 30.8 | 2.3 | * | * | 0.9 | 1.3 | 1.9 | 3.6 | 11.0 | 23.9 | 80.0 |
| 1999..... | 4.5 | 24.0 | 1.8 | * | * | 1.1 | 1.4 | 2.8 | 5.4 | 11.5 | 36.5 | 90.5 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Intentional self-harm (suicide) (X60-X84,Y87.0) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 10.7 | ... | ... | 0.8 | 10.4 | 12.8 | 14.6 | 14.6 | 12.3 | 12.6 | 17.7 | 19.4 |
| 1999..... | 10.7 | ... | ... | 0.6 | 10.3 | 13.5 | 14.4 | 14.2 | 12.4 | 13.6 | 18.3 | 19.2 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 17.5 | ... | ... | 1.2 | 17.4 | 21.2 | 22.9 | 22.8 | 19.8 | 23.0 | 38.0 | 54.5 |
| 1999..... | 17.6 | ... | ... | 1.0 | 17.2 | 22.3 | 22.6 | 22.0 | 20.2 | 25.0 | 38.3 | 54.9 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.1 | ... | ... | 0.3 | 3.0 | 4.5 | 6.4 | 6.8 | 5.4 | 4.1 | 4.0 | 4.3 |
| 1999..... | 4.1 | ... | ... | 0.3 | 3.1 | 4.8 | 6.4 | 6.7 | 5.2 | 4.2 | 4.8 | 4.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 11.7 | ... | ... | 0.8 | 10.9 | 13.7 | 16.1 | 16.1 | 13.5 | 13.7 | 18.8 | 20.8 |
| 1999..... | 11.7 | ... | ... | 0.7 | 10.7 | 14.3 | 15.9 | 15.6 | 13.4 | 14.6 | 19.4 | 20.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 19.1 | ... | ... | 1.3 | 18.2 | 22.4 | 25.0 | 24.8 | 21.5 | 24.7 | 40.4 | 58.8 |
| 1999..... | 19.1 | ... | ... | 1.0 | 17.8 | 23.4 | 24.5 | 23.9 | 21.6 | 26.8 | 40.8 | 59.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 4.5 | ... | ... | 0.4 | 3.1 | 4.8 | 7.2 | 7.6 | 6.0 | 4.4 | 4.1 | 4.5 |
| 1999..... | 4.5 | ... | ... | 0.3 | 3.2 | 5.1 | 7.2 | 7.5 | 5.7 | 4.5 | 5.0 | 4.3 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 5.6 | ... | ... | 0.6 | 8.2 | 9.1 | 7.7 | 6.7 | 4.3 | 5.0 | 5.9 | * |
| 1999..... | 5.6 | ... | ... | 0.5 | 8.2 | 9.5 | 7.7 | 5.9 | 5.0 | 5.7 | 5.8 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Intentional self-harm (suicide) (X60-X84,Y87.0)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|-----|-----|-----|------|------|------|------|-----|------|------|---|
| 2000..... | 9.8 | ... | ... | 1.1 | 14.2 | 16.2 | 13.3 | 11.7 | 7.9 | 10.6 | 11.8 | * |
| 1999..... | 10.0 | ... | ... | 0.7 | 14.4 | 17.4 | 13.4 | 10.5 | 9.6 | 11.3 | 13.5 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|-----|-----|-----|-----|-----|---|---|---|
| 2000..... | 1.8 | ... | ... | * | 2.2 | 2.7 | 2.7 | 2.5 | 1.5 | * | * | * |
| 1999..... | 1.6 | ... | ... | * | 2.0 | 2.4 | 2.6 | 2.0 | 1.6 | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

ALL RACES 3/, BOTH SEXES

| ALL RACES 3/, MALE | | | | | | | | | | | | |
|--------------------|------|-----|-----|-----|------|------|------|------|------|------|------|------|
| 2000..... | 10.7 | ... | ... | 0.4 | 10.4 | 11.4 | 11.9 | 13.3 | 13.7 | 18.6 | 30.3 | 40.7 |
| 1999..... | 10.9 | ... | ... | 0.4 | 10.8 | 12.1 | 11.6 | 12.9 | 14.3 | 19.7 | 30.8 | 40.0 |

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2000..... | 1.5 | ... | ... | 0.1 | 1.2 | 1.8 | 2.4 | 2.4 | 2.1 | 1.7 | 1.4 | 1.1 |
| 1999..... | 1.5 | ... | ... | 0.1 | 1.3 | 1.8 | 2.3 | 2.3 | 2.2 | 1.8 | 1.5 | 1.0 |

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| 2000..... | 6.7 | ... | ... | 0.3 | 6.2 | 7.0 | 8.0 | 8.6 | 8.4 | 10.2 | 14.2 | 14.1 |
| 1999..... | 6.7 | ... | ... | 0.3 | 6.3 | 7.4 | 7.7 | 8.3 | 8.7 | 10.7 | 14.2 | 13.6 |

| | | | | | | | | | | | | |
|-----------|------|-----|-----|-----|------|------|------|------|------|------|------|------|
| 2000..... | 11.9 | ... | ... | 0.5 | 10.8 | 12.1 | 13.1 | 14.6 | 14.9 | 20.1 | 32.8 | 44.3 |
| 1999..... | 11.9 | ... | ... | 0.5 | 11.1 | 12.7 | 12.7 | 14.1 | 15.4 | 21.3 | 32.9 | 43.5 |

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2000..... | 1.7 | ... | ... | * | 1.3 | 1.9 | 2.7 | 2.6 | 2.4 | 2.0 | 1.5 | 1.2 |
| 1999..... | 1.7 | ... | ... | 0.1 | 1.3 | 2.1 | 2.6 | 2.6 | 2.4 | 1.9 | 1.6 | 1.0 |

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 3.1 | ... | ... | * | 5.3 | 5.2 | 3.4 | 3.3 | 3.0 | 3.7 | 3.6 | * |
| 1999..... | 3.2 | ... | ... | * | 5.7 | 5.1 | 3.5 | 3.1 | 3.5 | 3.9 | 4.8 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|------|------|-----|-----|-----|-----|------|---|
| 2000..... | 5.8 | ... | ... | * | 9.6 | 9.7 | 6.3 | 6.1 | 5.9 | 8.4 | 7.9 | * |
| 1999..... | 6.1 | ... | ... | * | 10.3 | 10.0 | 6.7 | 5.8 | 7.3 | 8.3 | 11.7 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|-----|-----|-----|-----|---|---|---|---|
| 2000..... | 0.7 | ... | ... | * | 0.9 | 1.2 | 0.8 | 1.1 | * | * | * | * |
| 1999..... | 0.6 | ... | ... | * | 1.0 | 0.8 | 0.7 | * | * | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEARALL
AGES 1/
UNDER 1
YEAR 2/
1-4
YEARS
5-14
YEARS
15-24
YEARS
25-34
YEARS
35-44
YEARS
45-54
YEARS
55-64
YEARS
65-74
YEARS
75-84
YEARS
85 YEARS
AND OVER

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2000..... | 4.6 | ... | ... | 0.5 | 4.5 | 6.2 | 7.5 | 6.9 | 4.6 | 3.3 | 4.6 | 6.4 |
| 1999..... | 4.6 | ... | ... | 0.4 | 4.2 | 6.6 | 7.5 | 6.7 | 4.4 | 3.7 | 5.0 | 6.6 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|------|------|-----|-----|-----|-----|------|
| 2000..... | 6.8 | ... | ... | 0.8 | 7.0 | 9.8 | 11.0 | 9.6 | 6.1 | 4.4 | 7.6 | 13.8 |
| 1999..... | 6.7 | ... | ... | 0.6 | 6.4 | 10.2 | 10.9 | 9.2 | 5.9 | 5.3 | 7.6 | 14.9 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2000..... | 2.6 | ... | ... | 0.2 | 1.8 | 2.8 | 4.0 | 4.4 | 3.3 | 2.3 | 2.6 | 3.2 |
| 1999..... | 2.6 | ... | ... | 0.1 | 1.9 | 2.9 | 4.1 | 4.4 | 3.1 | 2.4 | 3.3 | 3.1 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2000..... | 5.0 | ... | ... | 0.5 | 4.7 | 6.6 | 8.2 | 7.5 | 5.1 | 3.4 | 4.7 | 6.7 |
| 1999..... | 5.0 | ... | ... | 0.4 | 4.4 | 6.9 | 8.2 | 7.3 | 4.7 | 3.9 | 5.2 | 7.1 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|------|------|------|-----|-----|-----|------|
| 2000..... | 7.3 | ... | ... | 0.7 | 7.4 | 10.3 | 11.8 | 10.2 | 6.6 | 4.5 | 7.7 | 14.5 |
| 1999..... | 7.2 | ... | ... | 0.6 | 6.8 | 10.7 | 11.8 | 9.8 | 6.2 | 5.5 | 7.9 | 16.1 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2000..... | 2.8 | ... | ... | 0.2 | 1.9 | 2.9 | 4.5 | 4.9 | 3.6 | 2.5 | 2.6 | 3.3 |
| 1999..... | 2.8 | ... | ... | 0.1 | 1.9 | 3.1 | 4.6 | 4.9 | 3.3 | 2.5 | 3.4 | 3.3 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 2.5 | ... | ... | 0.6 | 2.9 | 3.9 | 4.3 | 3.3 | 1.3 | 1.4 | 2.3 | * |
| 1999..... | 2.4 | ... | ... | 0.4 | 2.5 | 4.4 | 4.2 | 2.8 | 1.5 | 1.8 | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|---|
| 2000..... | 4.0 | ... | ... | 1.0 | 4.6 | 6.5 | 7.0 | 5.6 | 1.9 | * | * | * |
| 1999..... | 3.8 | ... | ... | * | 4.0 | 7.4 | 6.7 | 4.7 | 2.3 | 3.0 | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|-----|-----|-----|-----|---|---|---|---|
| 2000..... | 1.1 | ... | ... | * | 1.3 | 1.5 | 1.9 | 1.4 | * | * | * | * |
| 1999..... | 1.0 | ... | ... | * | 0.9 | 1.6 | 2.0 | 1.3 | * | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Assault (homicide) (X85-Y09,Y87.1) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.1 | 9.1 | 2.3 | 0.9 | 12.9 | 11.1 | 7.2 | 4.7 | 3.1 | 2.4 | 2.4 | 2.3 |
| 1999..... | 6.2 | 8.7 | 2.5 | 1.1 | 13.2 | 11.2 | 7.2 | 4.7 | 3.1 | 2.6 | 2.5 | 2.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 9.5 | 10.3 | 2.6 | 1.1 | 21.4 | 18.1 | 10.4 | 7.1 | 4.7 | 3.4 | 3.1 | 3.1 |
| 1999..... | 9.6 | 9.6 | 2.5 | 1.2 | 21.7 | 17.8 | 10.4 | 7.0 | 4.9 | 3.7 | 3.3 | 3.3 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.8 | 7.8 | 2.1 | 0.7 | 3.9 | 4.3 | 4.0 | 2.5 | 1.6 | 1.6 | 2.0 | 2.0 |
| 1999..... | 2.9 | 7.7 | 2.5 | 0.9 | 4.4 | 4.6 | 4.0 | 2.4 | 1.5 | 1.7 | 2.0 | 2.0 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.7 | 6.5 | 1.7 | 0.8 | 6.5 | 6.1 | 4.7 | 3.3 | 2.4 | 1.9 | 2.0 | 2.1 |
| 1999..... | 3.8 | 6.6 | 1.7 | 0.8 | 6.9 | 6.3 | 4.8 | 3.4 | 2.5 | 2.2 | 2.1 | 2.1 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 5.3 | 8.1 | 1.9 | 0.9 | 10.1 | 9.0 | 6.5 | 4.6 | 3.5 | 2.4 | 2.4 | 2.7 |
| 1999..... | 5.6 | 7.8 | 1.8 | 0.9 | 10.5 | 9.5 | 6.7 | 4.9 | 3.7 | 3.0 | 2.7 | 2.6 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 2.1 | 4.9 | 1.4 | 0.6 | 2.7 | 3.1 | 2.8 | 2.1 | 1.4 | 1.5 | 1.8 | 1.9 |
| 1999..... | 2.2 | 5.3 | 1.7 | 0.7 | 3.0 | 3.2 | 2.9 | 1.8 | 1.3 | 1.6 | 1.6 | 1.9 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 22.3 | 24.0 | 6.5 | 2.0 | 48.3 | 42.5 | 24.1 | 15.2 | 8.5 | 7.4 | 7.1 | * |
| 1999..... | 21.9 | 20.7 | 6.8 | 2.4 | 48.4 | 40.8 | 23.4 | 14.5 | 8.3 | 6.2 | 7.5 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Assault (homicide) (X85-Y09,Y87.1) | | | | | | | | | | | | |

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|------|-----|-----|------|------|------|------|------|------|------|---|
| 2000..... | 38.6 | 24.6 | 6.6 | 2.3 | 85.3 | 76.9 | 38.8 | 27.0 | 15.3 | 13.3 | 11.5 | * |
| 1999..... | 37.5 | 20.8 | 6.6 | 2.9 | 85.1 | 71.8 | 37.5 | 24.6 | 15.8 | 11.3 | 11.7 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|------|------|------|-----|-----|-----|-----|---|
| 2000..... | 7.5 | 23.4 | 6.4 | 1.7 | 11.0 | 11.4 | 11.1 | 5.3 | 3.2 | 3.1 | 4.5 | * |
| 1999..... | 7.8 | 20.7 | 7.0 | 1.8 | 11.5 | 12.8 | 11.0 | 6.1 | 2.6 | 2.5 | 5.0 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Assault (homicide) by discharge of firearms (X93-X95) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 3.9 | * | 0.2 | 0.5 | 10.3 | 8.2 | 4.2 | 2.6 | 1.5 | 1.0 | 0.7 | 0.7 |
| 1999..... | 4.0 | * | 0.3 | 0.6 | 10.7 | 8.0 | 4.2 | 2.5 | 1.6 | 1.1 | 0.8 | 0.6 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 6.7 | * | * | 0.7 | 18.1 | 14.2 | 6.7 | 4.0 | 2.3 | 1.3 | 0.7 | * |
| 1999..... | 6.7 | * | 0.3 | 0.8 | 18.5 | 13.9 | 6.4 | 3.9 | 2.6 | 1.5 | 1.0 | * |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.3 | * | * | 0.3 | 2.2 | 2.3 | 1.9 | 1.2 | 0.8 | 0.8 | 0.6 | * |
| 1999..... | 1.4 | * | 0.3 | 0.4 | 2.5 | 2.3 | 2.0 | 1.2 | 0.6 | 0.7 | 0.6 | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.1 | * | * | 0.4 | 4.7 | 4.0 | 2.7 | 1.8 | 1.3 | 0.8 | 0.6 | 0.6 |
| 1999..... | 2.2 | * | 0.2 | 0.4 | 4.9 | 4.1 | 2.7 | 1.7 | 1.3 | 0.9 | 0.7 | 0.6 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 3.3 | * | * | 0.5 | 7.8 | 6.4 | 3.9 | 2.6 | 1.8 | 1.0 | 0.6 | * |
| 1999..... | 3.5 | * | * | 0.5 | 8.1 | 6.6 | 4.0 | 2.6 | 1.9 | 1.2 | 0.7 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.0 | * | * | 0.2 | 1.4 | 1.6 | 1.4 | 1.0 | 0.8 | 0.7 | 0.6 | * |
| 1999..... | 1.0 | * | * | 0.3 | 1.6 | 1.5 | 1.5 | 0.9 | 0.6 | 0.7 | 0.6 | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 16.1 | * | * | 1.1 | 41.7 | 34.6 | 15.1 | 8.6 | 3.5 | 2.7 | * | * |
| 1999..... | 16.0 | * | * | 1.3 | 42.7 | 32.6 | 14.2 | 8.3 | 4.4 | 2.3 | 2.4 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Assault (homicide) by discharge of firearms (X93-X95)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|------|---|---|-----|------|------|------|------|-----|-----|---|---|
| 2000..... | 30.3 | * | * | 1.7 | 76.5 | 65.9 | 26.9 | 15.8 | 6.6 | 5.2 | * | * |
| 1999..... | 29.5 | * | * | 1.9 | 77.3 | 61.4 | 24.2 | 14.6 | 9.3 | 4.4 | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|-----|-----|-----|-----|-----|---|---|---|---|
| 2000..... | 3.3 | * | * | * | 6.6 | 6.4 | 4.6 | 2.6 | * | * | * | * |
| 1999..... | 3.8 | * | * | 0.7 | 7.8 | 6.7 | 5.4 | 3.1 | * | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 2.2 | 8.8 | 2.2 | 0.5 | 2.5 | 2.9 | 2.9 | 2.1 | 1.5 | 1.4 | 1.8 | 1.7 |
| 1999..... | 2.2 | 8.5 | 2.2 | 0.5 | 2.5 | 3.1 | 3.0 | 2.2 | 1.5 | 1.5 | 1.7 | 1.8 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 2.8 | 9.9 | 2.3 | 0.4 | 3.3 | 3.8 | 3.7 | 3.0 | 2.4 | 2.1 | 2.4 | 2.4 |
| 1999..... | 2.9 | 9.3 | 2.2 | 0.5 | 3.2 | 3.9 | 4.0 | 3.2 | 2.2 | 2.2 | 2.3 | 2.5 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.5 | 7.6 | 2.0 | 0.5 | 1.7 | 2.0 | 2.1 | 1.3 | 0.8 | 0.9 | 1.4 | 1.4 |
| 1999..... | 1.6 | 7.5 | 2.1 | 0.6 | 1.8 | 2.3 | 2.0 | 1.2 | 0.8 | 1.0 | 1.3 | 1.5 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.6 | 6.3 | 1.5 | 0.4 | 1.8 | 2.1 | 2.0 | 1.5 | 1.2 | 1.1 | 1.5 | 1.5 |
| 1999..... | 1.7 | 6.5 | 1.5 | 0.4 | 1.9 | 2.3 | 2.0 | 1.6 | 1.2 | 1.3 | 1.4 | 1.5 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 2.0 | 7.9 | 1.7 | 0.4 | 2.3 | 2.7 | 2.6 | 2.0 | 1.7 | 1.5 | 1.9 | 2.0 |
| 1999..... | 2.1 | 7.6 | 1.5 | 0.4 | 2.4 | 2.9 | 2.7 | 2.3 | 1.8 | 1.7 | 1.9 | 1.8 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 1.1 | 4.7 | 1.3 | 0.4 | 1.3 | 1.5 | 1.4 | 1.1 | 0.6 | 0.7 | 1.2 | 1.3 |
| 1999..... | 1.2 | 5.3 | 1.5 | 0.5 | 1.5 | 1.7 | 1.4 | 0.9 | 0.7 | 0.9 | 1.0 | 1.4 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 6.1 | 22.8 | 6.2 | 0.9 | 6.6 | 7.9 | 9.0 | 6.6 | 4.9 | 4.7 | 5.2 | * |
| 1999..... | 5.9 | 20.4 | 6.0 | 1.0 | 5.8 | 8.2 | 9.2 | 6.2 | 3.9 | 3.9 | 5.2 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTER- NATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|---|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|------|------|------|-----|-----|-----|---|
| 2000..... | 8.3 | 22.3 | 6.2 | * | 8.8 | 11.0 | 11.8 | 11.3 | 8.7 | 8.1 | 7.9 | * |
| 1999..... | 8.0 | 20.4 | 6.0 | 0.9 | 7.8 | 10.5 | 13.3 | 10.0 | 6.5 | 6.9 | 6.9 | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 2000..... | 4.2 | 23.4 | 6.1 | 1.2 | 4.4 | 5.0 | 6.5 | 2.8 | 2.0 | 2.2 | 3.6 | * |
| 1999..... | 4.0 | 20.4 | 6.0 | 1.1 | 3.7 | 6.1 | 5.6 | 3.0 | 1.9 | * | 4.1 | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--------------------------------|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Legal intervention (Y35,Y89.0) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | 0.2 | 0.3 | 0.2 | 0.1 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | 0.2 | 0.4 | 0.3 | 0.1 | 0.1 | * | * | * |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | 0.4 | 0.6 | 0.4 | 0.3 | * | * | * | * |
| 1999..... | 0.3 | * | * | * | 0.3 | 0.7 | 0.5 | 0.3 | 0.2 | * | * | * |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | * | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | 0.1 | 0.2 | 0.2 | 0.1 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | 0.1 | 0.3 | 0.2 | 0.1 | * | * | * | * |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | 0.3 | 0.5 | 0.3 | 0.3 | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.2 | 0.6 | 0.4 | 0.3 | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | * | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.3 | * | * | * | 0.5 | 0.6 | 0.6 | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | 0.6 | 0.8 | 0.6 | * | * | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|---|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Events of undetermined intent (Y10-Y34,Y87.2,Y89.9) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.4 | 1.7 | 0.3 | 0.2 | 1.0 | 1.8 | 2.9 | 2.3 | 1.0 | 0.6 | 0.6 | 1.3 |
| 1999..... | 1.4 | 1.4 | 0.3 | 0.1 | 1.1 | 2.0 | 2.9 | 2.2 | 0.9 | 0.6 | 0.8 | 1.4 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 2.0 | 1.4 | 0.3 | 0.2 | 1.5 | 2.6 | 4.0 | 3.1 | 1.3 | 0.9 | 1.0 | 1.6 |
| 1999..... | 2.0 | 1.7 | 0.3 | 0.2 | 1.8 | 2.8 | 4.0 | 3.0 | 1.3 | 0.9 | 1.1 | 2.0 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.8 | 2.0 | * | * | 0.5 | 1.0 | 1.8 | 1.4 | 0.6 | 0.3 | 0.4 | 1.2 |
| 1999..... | 0.9 | 1.2 | 0.3 | * | 0.5 | 1.2 | 1.8 | 1.4 | 0.6 | 0.4 | 0.6 | 1.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | 1.4 | 0.2 | 0.1 | 1.0 | 1.9 | 2.8 | 2.1 | 0.9 | 0.5 | 0.6 | 1.2 |
| 1999..... | 1.4 | 1.2 | 0.2 | 0.1 | 1.1 | 2.0 | 2.8 | 2.0 | 0.9 | 0.6 | 0.8 | 1.4 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | 0.2 | 1.5 | 2.6 | 3.8 | 2.9 | 1.1 | 0.7 | 1.0 | * |
| 1999..... | 1.9 | 1.4 | * | 0.2 | 1.8 | 2.7 | 3.8 | 2.6 | 1.2 | 0.8 | 1.1 | 2.1 |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.8 | 1.6 | * | * | 0.5 | 1.1 | 1.7 | 1.4 | 0.6 | 0.4 | 0.4 | 1.1 |
| 1999..... | 0.9 | * | * | * | 0.4 | 1.3 | 1.8 | 1.5 | 0.7 | 0.4 | 0.6 | 1.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.9 | * | * | 0.3 | 0.9 | 2.0 | 4.1 | 3.8 | 1.8 | * | * | * |
| 1999..... | 2.0 | * | * | * | 1.2 | 2.3 | 4.2 | 4.3 | 1.2 | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|---|---|---|
| 2000..... | 2.9 | * | * | * | 1.6 | 3.2 | 6.0 | 6.1 | 3.2 | * | * | * |
| 1999..... | 3.2 | * | * | * | 1.9 | 3.7 | 6.4 | 7.5 | 2.7 | * | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|---|---|---|---|
| 2000..... | 1.0 | * | * | * | * | 0.9 | 2.4 | 1.8 | * | * | * | * |
| 1999..... | 0.9 | * | * | * | * | 1.0 | 2.3 | 1.6 | * | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Discharge of firearms, undetermined intent (Y22-Y24) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | 0.2 | 0.1 | 0.1 | 0.1 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | 0.3 | 0.1 | 0.1 | 0.1 | * | * | * | * |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | 0.3 | 0.2 | 0.1 | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.6 | 0.2 | 0.2 | 0.1 | * | * | * | * |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | 0.2 | 0.1 | 0.1 | 0.1 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | 0.3 | 0.1 | 0.1 | 0.1 | * | * | * | * |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 0.1 | * | * | * | 0.3 | 0.2 | 0.1 | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.5 | 0.2 | 0.2 | * | * | * | * | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 0.2 | * | * | * | 0.3 | * | * | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.6 | * | * | * | * | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| | ALL AGES 1/ YEAR 2/ YEARS | UNDER 1 YEAR 2/ YEARS | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|------------------------------------|-----------------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9) | | | | | | | | | | | | |
| ALL RACES 3/, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | 1.7 | 0.3 | 0.1 | 0.8 | 1.7 | 2.8 | 2.2 | 0.9 | 0.5 | 0.6 | 1.3 |
| 1999..... | 1.3 | 1.4 | 0.3 | 0.1 | 0.8 | 1.8 | 2.8 | 2.1 | 0.9 | 0.5 | 0.7 | 1.3 |
| ALL RACES 3/, MALE | | | | | | | | | | | | |
| 2000..... | 1.8 | 1.4 | 0.3 | 0.2 | 1.1 | 2.4 | 3.8 | 3.0 | 1.2 | 0.7 | 0.9 | * |
| 1999..... | 1.8 | 1.7 | * | 0.1 | 1.2 | 2.5 | 3.8 | 2.9 | 1.2 | 0.7 | 0.9 | 1.7 |
| ALL RACES 3/, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.8 | 2.0 | * | * | 0.4 | 1.0 | 1.7 | 1.4 | 0.6 | 0.3 | 0.4 | 1.2 |
| 1999..... | 0.8 | 1.2 | 0.3 | * | 0.4 | 1.1 | 1.8 | 1.4 | 0.6 | 0.4 | 0.5 | 1.1 |
| WHITE, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.3 | 1.4 | 0.2 | 0.1 | 0.8 | 1.7 | 2.7 | 2.1 | 0.8 | 0.5 | 0.6 | 1.2 |
| 1999..... | 1.3 | 1.2 | 0.2 | 0.1 | 0.9 | 1.9 | 2.6 | 1.9 | 0.9 | 0.5 | 0.7 | 1.3 |
| WHITE, MALE | | | | | | | | | | | | |
| 2000..... | 1.7 | * | * | 0.2 | 1.2 | 2.4 | 3.7 | 2.8 | 1.1 | 0.6 | 0.9 | * |
| 1999..... | 1.7 | 1.4 | * | 0.1 | 1.3 | 2.5 | 3.5 | 2.5 | 1.1 | 0.7 | 0.9 | * |
| WHITE, FEMALE | | | | | | | | | | | | |
| 2000..... | 0.8 | 1.6 | * | * | 0.5 | 1.0 | 1.7 | 1.4 | 0.6 | 0.3 | 0.4 | 1.1 |
| 1999..... | 0.9 | * | * | * | 0.4 | 1.2 | 1.7 | 1.4 | 0.7 | 0.4 | 0.6 | 1.1 |
| BLACK, BOTH SEXES | | | | | | | | | | | | |
| 2000..... | 1.7 | * | * | * | 0.5 | 1.7 | 4.0 | 3.7 | 1.7 | * | * | * |
| 1999..... | 1.8 | * | * | * | 0.6 | 2.0 | 4.1 | 4.2 | 1.2 | * | * | * |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992),

| CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|-----|-----|-----|-----|-----|---|---|---|
| 2000..... | 2.6 | * | * | * | 1.0 | 2.7 | 5.9 | 6.0 | 2.9 | * | * | * |
| 1999..... | 2.8 | * | * | * | 0.9 | 3.2 | 6.3 | 7.3 | 2.6 | * | * | * |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|---|---|---|---|
| 2000..... | 0.9 | * | * | * | * | 0.8 | 2.3 | 1.8 | * | * | * | * |
| 1999..... | 0.9 | * | * | * | * | 0.9 | 2.2 | 1.6 | * | * | * | * |

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| ALL AGES 1/ RACE, SEX, AND YEAR | UNDER 1 YEAR 2/ AGES 1/ RACE, SEX, AND YEAR | 1-4 YEARS AGES 1/ RACE, SEX, AND YEAR | 5-14 YEARS AGES 1/ RACE, SEX, AND YEAR | 15-24 YEARS AGES 1/ RACE, SEX, AND YEAR | 25-34 YEARS AGES 1/ RACE, SEX, AND YEAR | 35-44 YEARS AGES 1/ RACE, SEX, AND YEAR | 45-54 YEARS AGES 1/ RACE, SEX, AND YEAR | 55-64 YEARS AGES 1/ RACE, SEX, AND YEAR | 65-74 YEARS AGES 1/ RACE, SEX, AND YEAR | 75-84 YEARS AGES 1/ RACE, SEX, AND YEAR | 85 YEARS AND OVER AGES 1/ RACE, SEX, AND YEAR |
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|
|---------------------------------------|--|--|---|--|--|--|--|--|--|--|--|

ALL RACES 3/, BOTH SEXES

ALL RACES 3/, MALE

ALL RACES 3/, FEMALE

WHITE, BOTH SEXES

WHITE, MALE

WHITE, FEMALE

BLACK, BOTH SEXES

[illegible]

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

| OF DISEASES, 1992), RACE, SEX, AND YEAR | ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

BLACK, MALE

BLACK, FEMALE

[illegible]

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEARALL
AGES 1/
UNDER 1
YEAR 2/
1-4
YEARS
5-14
YEARS
15-24
YEARS
25-34
YEARS
35-44
YEARS
45-54
YEARS
55-64
YEARS
65-74
YEARS
75-84
YEARS
85 YEARS
AND OVER

Complications of medical and surgical care (Y40-Y84,Y88)

ALL RACES 3/, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| 2000..... | 1.1 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 1.6 | 3.6 | 6.7 | 13.5 |
| 1999..... | 1.0 | 1.4 | 0.2 | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 3.2 | 6.7 | 12.7 |

ALL RACES 3/, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|---|---|-----|-----|-----|-----|-----|-----|-----|------|
| 2000..... | 1.1 | * | * | * | * | 0.3 | 0.3 | 0.8 | 1.6 | 4.0 | 8.1 | 15.2 |
| 1999..... | 1.0 | 1.6 | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.5 | 3.4 | 7.7 | 15.9 |

ALL RACES 3/, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|-----|---|---|---|-----|-----|-----|-----|-----|-----|------|
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.6 | 3.2 | 5.8 | 12.7 |
| 1999..... | 1.1 | 1.2 | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 3.1 | 6.0 | 11.3 |

WHITE, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| 2000..... | 1.1 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.7 | 1.4 | 3.4 | 6.5 | 13.0 |
| 1999..... | 1.0 | 1.2 | 0.2 | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.2 | 3.1 | 6.4 | 12.1 |

WHITE, MALE

| | | | | | | | | | | | | |
|-----------|-----|-----|---|---|---|-----|-----|-----|-----|-----|-----|------|
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.4 | 3.8 | 8.1 | 15.1 |
| 1999..... | 1.0 | 1.4 | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 3.2 | 7.4 | 15.1 |

WHITE, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|-----|-----|------|
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.3 | 3.0 | 5.4 | 12.1 |
| 1999..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.9 | 5.7 | 10.8 |

BLACK, BOTH SEXES

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|-----|-----|-----|-----|-----|------|------|
| 2000..... | 1.4 | * | * | * | * | 0.4 | 0.5 | 1.4 | 3.6 | 5.9 | 10.7 | 21.0 |
| 1999..... | 1.3 | * | * | * | * | * | 0.6 | 1.3 | 2.6 | 5.1 | 10.8 | 20.1 |

TABLE 290. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1999-2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, SEX, AND YEAR

| ALL AGES 1/ | UNDER 1 YEAR 2/ | 1-4 YEARS | 5-14 YEARS | 15-24 YEARS | 25-34 YEARS | 35-44 YEARS | 45-54 YEARS | 55-64 YEARS | 65-74 YEARS | 75-84 YEARS | 85 YEARS AND OVER |
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|----------------|--------------------|--------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Complications of medical and surgical care (Y40-Y84,Y88)

BLACK, MALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|------|------|
| 2000..... | 1.2 | * | * | * | * | * | * | 1.3 | 3.3 | 7.3 | 10.0 | * |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.5 | 3.0 | 5.7 | 12.0 | 21.7 |

BLACK, FEMALE

| | | | | | | | | | | | | |
|-----------|-----|---|---|---|---|---|---|-----|-----|-----|------|------|
| 2000..... | 1.6 | * | * | * | * | * | * | 1.5 | 3.8 | 5.0 | 11.1 | 22.7 |
| 1999..... | 1.4 | * | * | * | * | * | * | 1.2 | 2.4 | 4.7 | 10.1 | 19.5 |

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.